

ON THE σ -FINITENESS OF A VARIATIONAL MEASURE

DIANA CAPONETTI, Arcavacata di Rende

(Received November 23, 2001)

Abstract. The σ -finiteness of a variational measure, generated by a real valued function, is proved whenever it is σ -finite on all Borel sets that are negligible with respect to a σ -finite variational measure generated by a continuous function.

Keywords: variational measure, H -differentiable, H -density

MSC 2000: 26A39, 26A24

1. INTRODUCTION

In 1994, a question was posed by W. Pfeffer (see [13]) whether the absolute continuity of a variational measure, generated by a real valued function, with respect to the Lebesgue measure would imply its σ -finiteness. The affirmative answer was first given in [2], providing a full descriptive characterization of the Henstock-Kurzweil integral (see also [14], and [4], [5], [6], [8] for higher dimensional results). Then in [18], strengthening the result presented in [2], the author proved that a variational measure is σ -finite whenever it is σ -finite on all subsets of zero Lebesgue measure (see also [3] for a variational measure related to a certain class of differentiation bases). In this paper we show that the same result holds if the Lebesgue measure is replaced by a suitable variational measure. Namely, the variational measure V_*F , generated by a function $F: [a, b] \rightarrow \mathbb{R}$, is σ -finite on $[a, b]$ whenever it is σ -finite on all subsets having measure zero with respect to a σ -finite variational measure V_*U generated by a continuous function $U: [a, b] \rightarrow \mathbb{R}$. We derive some results on the differentiability of the function F with respect to U , and a representation theorem for the variational measure V_*F in terms of the Lebesgue integral.

Supported by M.I.U.R. of Italy.

2. PRELIMINARIES

If $E \subset \mathbb{R}$, then $|E|$ and $\text{int}E$ denote the outer Lebesgue measure and the interior of E , respectively. All functions we consider are real-valued. By $(\mathcal{L}) \int$ we denote the Lebesgue integral. We always consider nondegenerate subintervals of \mathbb{R} . For $c, d \in \mathbb{R}$ with $c < d$, we denote by $[c, d]$ the compact subinterval of \mathbb{R} with endpoints c and d , and by (c, d) the open one. A collection of intervals is called *nonoverlapping* whenever their interiors are disjoint. Throughout this note $[a, b]$ will be a fixed interval. A *partition in* $[a, b]$ is a collection $P = \{([a_1, b_1], x_1), \dots, ([a_p, b_p], x_p)\}$ where $[a_1, b_1], \dots, [a_p, b_p]$ are nonoverlapping subintervals of $[a, b]$ and $x_i \in [a_i, b_i]$ for $i = 1, \dots, p$. A positive function δ on $E \subset [a, b]$ is called a *gauge* on E . Given a gauge δ on $[a, b]$, a *partition* $P = \{([a_1, b_1], x_1), \dots, ([a_p, b_p], x_p)\}$ in $[a, b]$ is called

- (i) δ -*fine* if $b_i - a_i < \delta(x_i)$, $i = 1, \dots, p$;
- (ii) *of* $[a, b]$ if $\bigcup_{i=1}^p [a_i, b_i] = [a, b]$;
- (iii) *anchored in* E if $x_i \in E \subset [a, b]$ for each $i = 1, \dots, p$.

Let $H: [a, b] \rightarrow \mathbb{R}$ be a given function. The *variational measure* of H (see [17] and [2]) is the metric outer measure defined for each $E \subset [a, b]$ by

$$V_*H(E) = \inf_{\delta} \sup_P \sum_{i=1}^p |H(b_i) - H(a_i)|$$

where the infimum is taken over all gauges δ on E , and the supremum over all δ -fine partitions $P = \{([a_1, b_1], x_1), \dots, ([a_p, b_p], x_p)\}$ anchored in E .

If $V_*H(N) = 0$, then the set $N \subset [a, b]$ is called *H -negligible*. For details on metric outer measure we refer to [15] and [17]. We recall that H -negligible sets are V_*H -measurable, and any set that differs from a V_*H -measurable one by an H -negligible set is itself V_*H -measurable. We also recall that the restriction of a metric outer measure to the Borel sets is a measure.

V_*H is said to be σ -*finite* on $E \subset [a, b]$ if the set E is the union of sets E_n , $n = 1, 2, \dots$, satisfying $V_*H(E_n) < \infty$. A variational measure V_*F is said to be *absolutely continuous* with respect to V_*H if $V_*F(N) = 0$ for any H -negligible set $N \subset [a, b]$.

Remark 2.1. (i) Let $x \in [a, b]$. Then H is continuous at x if and only if $V_*H(\{x\}) = 0$.

(ii) If H is a continuous monotone function, then V_*H is the Lebesgue-Stieltjes measure associated with H , in which case

- (a) $V_*H([c, d]) = H(d) - H(c)$ for any subinterval $[c, d] \subset [a, b]$;

(b) V_*H is G_δ -regular, i.e. for every $E \subset [a, b]$ there is a V_*H -measurable G_δ set $Y \subset [a, b]$ containing E for which $V_*H(E) = V_*H(Y)$ (see [17, p. 62]).

According to [10, p. 416] a set $E \subset [a, b]$ is said to be H -null if it is the union of a countable set and an H -negligible set. A property is said to hold H -almost everywhere (abbreviated as H -a.e.) if the set of points where it fails to hold is H -null. However, if H is a continuous function, by Remark 2.1(i) we have that a set is H -null if and only if it is H -negligible.

Let F and H be any two functions on $[a, b]$. We need some definitions and results on the differentiability of the function F with respect to H . The *lower* and *upper derivative* of F with respect to H ,

$$\underline{D}_H F(x) = \liminf_{y \rightarrow x} \frac{F(y) - F(x)}{H(y) - H(x)} \quad \text{and} \quad \overline{D}_H F(x) = \limsup_{y \rightarrow x} \frac{F(y) - F(x)}{H(y) - H(x)},$$

are defined for all $x \in [a, b]$ for which $H(y) \neq H(x)$ in a neighborhood of x .

If $\underline{D}_H F(x) = \overline{D}_H F(x) \neq \pm\infty$ this common value is denoted by F'_H and F is said to be H -differentiable at x . Moreover, set

$$|\overline{D}|_H F(x) = \limsup_{y \rightarrow x} \frac{|F(y) - F(x)|}{|H(y) - H(x)|}.$$

The following result on H -differentiability will be useful. We point out that in [10] a function F is said to be VBG^o if V_*F is σ -finite on $[a, b]$.

Lemma 2.2 [10, Proposition 3.10]. *Let $F, H: [a, b] \rightarrow \mathbb{R}$ be given. If the variational measures V_*F and V_*H are σ -finite on $[a, b]$, then F is H -differentiable H -a.e. in $[a, b]$.*

The following lemma can be proved by standard arguments (cf. for example [12, Proposition 5.3.3]).

Lemma 2.3. *Let $F: [a, b] \rightarrow \mathbb{R}$ be given. If $H: [a, b] \rightarrow \mathbb{R}$ is a strictly increasing function, then for each $x \in [a, b]$ we have*

$$(1) \quad \overline{D}_H F(x) = \inf_{\delta} \sup_{[c, d]} \frac{F(d) - F(c)}{H(d) - H(c)}$$

where δ is a positive number and the supremum is taken over all subintervals $[c, d]$ of $[a, b]$ with $x \in [c, d]$ and $d - c < \delta$. If in addition H and F are continuous at x , then the supremum in (1) can be taken over all subintervals $[c, d]$ of $[a, b]$ with $x \in (c, d)$ and $d - c < \delta$.

Lemma 2.4. *Let $F: [a, b] \rightarrow \mathbb{R}$ be a continuous function. If $H: [a, b] \rightarrow \mathbb{R}$ is a continuous strictly increasing function, then $\overline{D}_H F$ is Borel-measurable.*

Proof. In view of Lemma 2.3, $\overline{D}_H F(x)$ can be written as in (1) where the supremum is taken over all subintervals $[c, d]$ of $[a, b]$ with $x \in (c, d)$ and $d - c < \delta$. Then by standard arguments (see for example [17, Theorem 4.2]), the upper derivative $\overline{D}_H F$ is Borel-measurable. \square

Clearly the same considerations of Lemma 2.3 and Lemma 2.4 apply to $\underline{D}_H F(x)$ and $|\overline{D}|_H F(x)$.

3. THE VARIATIONAL MEASURE

In order to study the properties of a variational measure, we introduce the following notion of H -density.

Definition 3.1. Let $H: [a, b] \rightarrow \mathbb{R}$ and let E be a subset of $[a, b]$. We say that a point $x \in [a, b]$ is a *point of H -density* for E if

$$\lim_{r \rightarrow 0^+} \frac{V_* H(E \cap [x - r, x + r])}{V_* H([x - r, x + r])} = 1.$$

The following lemma is a particular case of [11, Corollary 2.14].

Lemma 3.2. *Let $H: [a, b] \rightarrow \mathbb{R}$ be a continuous and strictly increasing function. Let E be a $V_* H$ -measurable subset of $[a, b]$. Then H -almost all points of E are H -density points for E .*

In view of Remark 2.1 (ii) we have that if $H: [a, b] \rightarrow \mathbb{R}$ is a continuous and strictly increasing function, then $V_* H$ is the corresponding Lebesgue-Stieltjes measure. Now we point out (see for example [7]) that the Vitali covering theorem holds for $V_* H$. Precisely, if a class of closed intervals covers a subset $A \subset [a, b]$ in the sense of Vitali, then there is a countable disjoint sequence of those intervals whose union differs from A by at most an H -negligible subset. In the following proposition we prove a result on the σ -finiteness of a variational measure by a technique similar to that used in [3, Theorem 3.1].

Proposition 3.3. *Let $F: [a, b] \rightarrow \mathbb{R}$ be given and let $H: [a, b] \rightarrow \mathbb{R}$ be a continuous and strictly increasing function. If $V_* F$ is σ -finite on all H -negligible Borel subsets of $[a, b]$, then $V_* F$ is σ -finite on $[a, b]$.*

Proof. Let Q be the set of all points $x \in [a, b]$ for which V_*F is not σ -finite on any open interval (c, d) of $[a, b]$ containing x . Clearly Q is closed and has no isolated points. Thus Q is a perfect set.

Now for any given interval $I \subset [a, b]$, let $\{I_j\}$ denote the sequence of intervals complementary to Q in I . Then a compactness argument shows that V_*F is σ -finite on I_j for each j . In particular, V_*F is σ -finite on the complement of Q in $[a, b]$. Therefore if $V_*H(Q) = 0$, by the hypothesis it follows that V_*F is σ -finite on $[a, b]$.

Assume by contradiction that $V_*H(Q) > 0$ and let K_Q be the set of all points of Q which are H -density points for Q . By Lemma 3.2, $V_*H(Q \setminus K_Q) = 0$. Let K denote the set of all $x \in K_Q$ for which the following condition holds: if $I \subset [a, b]$ is any interval containing x , then $V_*H(K_Q \cap \text{int}I) > 0$. We claim that $V_*H(K_Q \setminus K) = 0$. The family \mathcal{B} of all intervals $I \subset [a, b]$ for which $V_*H(K_Q \cap \text{int}I) = 0$ is a Vitali cover of the set $K_Q \setminus K$. By the Vitali covering theorem for Lebesgue-Stieltjes measures there is a disjoint sequence $\{I_{x_i}\}$ in \mathcal{B} with $x_i \in (K_Q \setminus K) \cap I_{x_i}$, such that

$$(2) \quad V_*H\left((K_Q \setminus K) \setminus \left(\bigcup_i I_{x_i}\right)\right) = 0.$$

For each i we have $V_*H(K_Q \cap \text{int}I_{x_i}) = 0$, which together with the continuity of H implies $V_*H(K_Q \cap I_{x_i}) = 0$. Then we have

$$(3) \quad V_*H\left(K_Q \cap \left(\bigcup_i I_{x_i}\right)\right) = 0.$$

Thus by (2) and (3) we have

$$V_*H(K_Q \setminus K) = V_*H\left((K_Q \setminus K) \setminus \left(\bigcup_i I_{x_i}\right)\right) + V_*H\left((K_Q \setminus K) \cap \left(\bigcup_i I_{x_i}\right)\right) = 0.$$

We show now that V_*F is not σ -finite on $K \cap I$, whenever I is an interval of $[a, b]$ which intersects K . As before let $\{I_j\}$ denote the sequence of intervals complementary to Q in I . Write

$$I = (K \cap I) \cup ((Q \setminus K) \cap I) \cup \left(\bigcup_j I_j\right),$$

and by Remark 2.1 (ii)(b) find an H -negligible G_δ set $Y \subset [a, b]$ containing $Q \setminus K$. Then we get

$$V_*F(I) \leq V_*F(K \cap I) + V_*F(Y \cap I) + V_*F\left(\bigcup_j I_j\right).$$

By the hypothesis V_*F is σ -finite on $Y \cap I$, and we have shown that it is σ -finite on $\bigcup_j I_j$. Hence the σ -finiteness of V_*F on $K \cap I$ would imply its σ -finiteness on I , which is not the case. This implies that for any gauge δ we have

$$(4) \quad \sup_P \sum_{i=1}^p |F(b_i) - F(a_i)| = \infty$$

where $P = \{([a_1, b_1], x_1), \dots, ([a_p, b_p], x_p)\}$ runs over all δ -fine partitions anchored in $K \cap I$.

Fix an open interval (c, d) containing a point of K . In view of Remark 2.1 (ii)(a), we may assume that $V_*H((c, d)) < 1/2$. Using (4) we can choose a finite collection $\{[a_i^{(1)}, b_i^{(1)}], i = 1, \dots, p_1\}$ of intervals contained in (c, d) , such that

$$\sum_{i=1}^{p_1} |F(b_i^{(1)}) - F(a_i^{(1)})| > 2.$$

We may assume that the family consists of at least two intervals. Also we have that the interior of each $[a_i^{(1)}, b_i^{(1)}]$ intersects K . Clearly,

$$\sum_{i=1}^{p_1} V_*H([a_i^{(1)}, b_i^{(1)}]) < 1/2.$$

Thinking of $[a, b]$ as $[a_1^{(0)}, b_1^{(0)}]$, we construct inductively finite collections $\{[a_i^{(k)}, b_i^{(k)}], i = 1, \dots, p_k\}$ such that the following conditions are satisfied for $k = 1, 2, \dots$:

- (i) $K \cap (a_i^{(k)}, b_i^{(k)}) \neq \emptyset$ for $i = 1, \dots, p_k$;
- (ii) each $[a_i^{(k)}, b_i^{(k)}]$ is contained in some $[a_j^{(k-1)}, b_j^{(k-1)}]$;
- (iii) each $[a_j^{(k-1)}, b_j^{(k-1)}]$ contains at least two intervals $[a_i^{(k)}, b_i^{(k)}]$;
- (iv) $\sum_{i=1}^{p_k} V_*H([a_i^{(k)}, b_i^{(k)}]) < 2^{-k}$;
- (v) $\sum_{i: [a_i^{(k)}, b_i^{(k)}] \subset [a_j^{(k-1)}, b_j^{(k-1)}]} |F(b_i^{(k)}) - F(a_i^{(k)})| > 2^k$ for each $j = 1, \dots, p_{k-1}$.

Now we define $N = \bigcap_{k=1}^{\infty} \bigcup_{i=1}^{p_k} [a_i^{(k)}, b_i^{(k)}]$. From conditions (i)–(iv) it follows that N is a perfect H -negligible set. As V_*F is σ -finite on N , we can write $N = \bigcup_{s=1}^{\infty} N_s$, where N_s are disjoint V_*F -measurable subsets of finite V_*F -measure. Choose a gauge δ on N such that for every integer $s \geq 1$

$$\sup_P \sum_{i=1}^p |F(b_i) - F(a_i)| < \infty$$

where $P = \{([a_1, b_1], x_1), \dots, ([a_p, b_p], x_p)\}$ runs over all δ -fine partitions anchored in N_s . Let $L_m = \{x \in N : \delta(x) > 1/m\}$ for $m = 1, 2, \dots$. Since $N = \bigcup_{m,s} (L_m \cap N_s)$, using the Baire category theorem we conclude that there exist integers m and s and an interval I with $N \cap I \neq \emptyset$ such that $L_m \cap N_s$ is a dense subset of $N \cap I$. We may assume $|I| < 1/m$. By the choice of δ we have

$$(5) \quad \sup_P \sum_{i=1}^p |F(b_i) - F(a_i)| < \infty$$

where $P = \{([a_1, b_1], x_1), \dots, ([a_p, b_p], x_p)\}$ runs over all δ -fine partitions anchored in $L_m \cap N_s$. Since I intersects N , then for all sufficiently large k there is some j such that $[a_j^{(k-1)}, b_j^{(k-1)}] \subset I$. Each interval $[a_i^{(k)}, b_i^{(k)}] \subset [a_j^{(k-1)}, b_j^{(k-1)}]$ contains a point of N and consequently a point, say x_{ik} , of $L_m \cap N_s$. Then $\{([a_i^{(k)}, b_i^{(k)}], x_{ik}) : [a_i^{(k)}, b_i^{(k)}] \subset [a_j^{(k-1)}, b_j^{(k-1)}]\}$ is a δ -fine partition anchored in $L_m \cap N_p$. Condition (v) implies

$$\sum_{i: [a_i^{(k)}, b_i^{(k)}] \subset [a_j^{(k-1)}, b_j^{(k-1)}]} |F(b_i^{(k)}) - F(a_i^{(k)})| > 2^k.$$

For a sufficiently large k , the last inequality contradicts (5), and the proposition is proved. \square

Theorem 3.4. *Let $F: [a, b] \rightarrow \mathbb{R}$ be given and let $U: [a, b] \rightarrow \mathbb{R}$ be a continuous function such that V_*U is σ -finite on $[a, b]$. If V_*F is σ -finite on all U -negligible Borel subsets of $[a, b]$, then V_*F is σ -finite on $[a, b]$.*

Proof. Since U is continuous we observe that V_*U coincides with the full variational measure ΔU^* introduced by Thomson in [17]. Then by [17, Theorem 7.8] the function U is VBG_* in the sense of Saks and by a theorem of Ward (see [16, p. 237]) there exists a continuous strictly increasing function H such that $|\overline{D}|_H U(x)$ is finite at every $x \in [a, b]$. Therefore by [10, Lemma 3.8], V_*U is absolutely continuous with respect to V_*H . This last property and the hypothesis imply that V_*F is σ -finite on all H -negligible Borel subsets of $[a, b]$. By Proposition 3.3, the σ -finiteness of V_*F on $[a, b]$ follows. \square

Corollary 3.5. *Let $F: [a, b] \rightarrow \mathbb{R}$ be given and let $U: [a, b] \rightarrow \mathbb{R}$ be a continuous function such that V_*U is σ -finite on $[a, b]$. If V_*F is σ -finite on all U -negligible Borel subsets of $[a, b]$, then F is U -differentiable U -a.e. in $[a, b]$.*

Proof. By Theorem 3.4, V_*F is σ -finite on $[a, b]$. Then the corollary follows from Lemma 2.2. \square

As a corollary of Theorem 3.4, we obtain a recently published result of V. Ene [9, Theorem 3.2]. We wish to point out that this result allows one to furnish a full descriptive characterization of the Henstock-Stieltjes integral introduced by Faure in [10] (see [9, Theorem 5.1 (iii)]).

Corollary 3.6. *Let $F: [a, b] \rightarrow \mathbb{R}$ be given and let $U: [a, b] \rightarrow \mathbb{R}$ be a continuous function such that V_*U is σ -finite on $[a, b]$. If V_*F is absolutely continuous with respect to V_*U , then V_*F is σ -finite on $[a, b]$.*

The following proposition allows us to represent V_*F on Borel sets in terms of the Lebesgue integral with respect to a σ -finite variational measure. It is based on a result of B. Bongiorno [1, Theorem 1] where a finite measure is considered.

Proposition 3.7. *Let $F: [a, b] \rightarrow \mathbb{R}$ be given and let $U: [a, b] \rightarrow \mathbb{R}$ be a continuous function such that V_*U is σ -finite on $[a, b]$. If V_*F is absolutely continuous with respect to V_*U , then*

$$(6) \quad V_*F(E) = (\mathcal{L}) \int_E |F'_U| dV_*U$$

for every Borel set $E \subset [a, b]$.

Proof. In view of Corollary 3.5 the variational measure V_*F is σ -finite on $[a, b]$. Therefore by Lemma 2.2, F'_U exists U -a.e. We observe that by the absolute continuity of V_*F with respect to V_*U and Remark 2.1(i), the function F is continuous. Let $E \subset [a, b]$ be a Borel set.

Assume first that U is strictly increasing. Since the set of all $x \in [a, b]$ for which $F'_U(x) \neq \overline{D}_U F(x)$ is U -negligible and by Lemma 2.4 $\overline{D}_U F$ is Borel-measurable, we have that F'_U is V_*U -measurable. Thus the Lebesgue integral $(\mathcal{L}) \int_E |F'_U| dV_*U$ exists (possibly equal to $+\infty$). By Remark 2.1(ii), V_*U is the Lebesgue-Stieltjes measure generated by U and $V_*U([c, d]) = U(d) - U(c)$. Thus F'_U coincides with the derivative of the set function $[c, d] \rightarrow F(d) - F(c)$ with respect to the measure V_*U .

Hence (6) follows by [1, Theorem 1] (cf. also [14, Proposition 10]).

Assume now V_*U to be σ -finite and let H denote, as in the proof of Theorem 3.3, a continuous strictly increasing function on $[a, b]$ such that V_*U is absolutely continuous with respect to V_*H . Then by the first part of the proof we get

$$(7) \quad V_*U(E) = (\mathcal{L}) \int_E |U'_H| dV_*H.$$

The hypothesis implies that V_*F is absolutely continuous with respect to V_*H , hence we also have

$$(8) \quad V_*F(E) = (\mathcal{L}) \int_E |F'_H| dV_*H.$$

Let N_1 denote the H -negligible, and hence U -negligible, subset of $[a, b]$ such that F'_H and U'_H exist for each $x \in [a, b] \setminus N_1$. Now let $N_2 = \{x \in [a, b] \setminus N_1 : U'_H(x) = 0\}$. We observe that N_2 is V_*H -measurable. Choose an $\varepsilon > 0$. Given $x \in N_2$, find a $\delta(x) > 0$ such that

$$|U(d) - U(c)| < \varepsilon(H(d) - H(c))$$

for any subinterval $[c, d]$ of $[a, b]$ with $x \in [c, d]$ and $d - c < \delta$. If $P = \{([a_1, b_1], x_1), \dots, ([a_p, b_p], x_p)\}$ is a δ -fine partition anchored in N_2 , then

$$\sum_{i=1}^p |U(b_i) - U(a_i)| < \varepsilon(H(b) - H(a)).$$

As ε is arbitrary, the set N_2 is U -negligible. Then the set $N = N_1 \cup N_2$ is U -negligible, and for any $x \in [a, b] \setminus N$ we have

$$(9) \quad F'_U(x) = F'_H(x)(U'_H(x))^{-1}.$$

Since by (7), for every V_*H -measurable function $g: [a, b] \rightarrow [0, \infty]$ we have

$$(\mathcal{L}) \int_E g \, dV_*U = (\mathcal{L}) \int_E |U'_H| g \, dV_*H,$$

by virtue of (8) and (9) the theorem follows for $g = |F'_U|$. □

A c k n o w l e d g m e n t. The author wish to thank Professor L. Di Piazza for her advice during the preparation of this paper.

References

- [1] *B. Bongiorno*: Essential variations. Springer Lecture Notes Math. 945 (1981), 187–193.
- [2] *B. Bongiorno, L. Di Piazza, V. Skvortsov*: A new full descriptive characterization of Denjoy-Perron integral. Real Anal. Exch. 21 (1995/96), 656–663.
- [3] *B. Bongiorno, L. Di Piazza, V. Skvortsov*: On variational measures related to some bases. J. Math. Anal. Appl. 250 (2000), 533–547.
- [4] *B. Bongiorno, L. Di Piazza, D. Preiss*: Infinite variation and derivatives in \mathbb{R}^m . J. Math. Anal. Appl. 224 (1998), 22–33.
- [5] *Z. Buczolich, W. F. Pfeffer*: When absolutely continuous implies σ -finite. Bull. Csi., Acad. Royale Belgique, serie 6 (1997), 155–160.
- [6] *Z. Buczolich, W. F. Pfeffer*: Variations of additive functions. Czechoslovak Math. J. 47 (1997), 525–555.
- [7] *J. L. Doob*: Measure Theory. Springer, New-York, 1994.
- [8] *L. Di Piazza*: Variational measures in the theory of the integration in \mathbb{R}^m . Czechoslovak Math. J. 51 (2001), 95–110.
- [9] *V. Ene*: Thomson's variational measure and nonabsolutely convergent integrals. Real Anal. Exch. 26 (2000/01), 35–50.

- [10] *C.-A. Faure*: A descriptive definition of the KH-Stieltjes integral. *Real Anal. Exch.* 23 (1997/98), 113–124.
- [11] *P. Mattila*: *Geometry of sets and measures in Euclidean spaces*. Cambridge University Press, 1995.
- [12] *W. F. Pfeffer*: *The Riemann Approach to Integration*. Cambridge University Press, 1993.
- [13] *W. F. Pfeffer*: On additive continuous functions of figures. *Rend. Istit. Mat. Univ. Trieste*, suppl. (1998), 115–133.
- [14] *W. F. Pfeffer*: The Lebesgue and Denjoy-Perron integrals from a descriptive point of view. *Ricerche Mat.* 48 (1999), 211–223.
- [15] *C. A. Rogers*: *Hausdorff Measures*. Cambridge, 1970.
- [16] *S. Saks*: *Theory of the Integral*. Dover, New York, 1964.
- [17] *B. S. Thomson*: Derivates of interval functions. *Mem. Amer. Math. Soc.*, Providence 452 (1991).
- [18] *B. S. Thomson*: Some properties of variational measures. *Real Anal. Exch.* 24 (1998/99), 845–853.

Author's address: Diana Caponetti, Dipartimento di Matematica, Università della Calabria, I-87036 Arcavacata di Rende (CS), Italy, e-mail: caponetti@unical.it.