

## OPERATIONAL QUANTITIES DERIVED FROM THE MINIMUM MODULUS

Manuel González<sup>†</sup> and Antonio Martín<sup>‡</sup>

**Abstract.** The minimum modulus  $\gamma(T)$  of an operator  $T$  is useful in perturbation theory because it characterizes the operators with closed range. Here we study the operational quantities derived from  $\gamma(T)$ . We show that the behavior of some of these quantities depends largely on whether the null space of  $T$  is finite dimensional or infinite dimensional.

### 1. Introduction

For every (linear bounded) non-zero operator  $T \in L(X, Y)$ , where  $X$  and  $Y$  are Banach spaces, the *minimum modulus* is defined by

$$\gamma(T) := \inf_{x \notin N(T)} \frac{\|Tx\|}{\text{dist}(x, N(T))}.$$

For  $T = 0$  we set  $\gamma(0) = 0$ . It is well known that  $\gamma(T) > 0$  if and only if  $T$  has closed range and  $T \neq 0$  [1, Theorem IV.1.6].

Here we study the operational quantities that can be derived from the minimum modulus  $\gamma(T)$ .

In the preliminaries, we give a description of the procedure to derive the quantities associated to a given quantity. This procedure, applied to the norm  $n(T) \equiv \|T\|$ , provides three quantities *in*, *sin* and *i\*n* which have been studied (with a different notation) in [9], [4], [7] and [2, 3]. Applied to the injection modulus  $j(T) := \inf\{\|Tx\| : x \in X, \|x\| = 1\}$ , it provides three quantities *s\*j*, *sj* and *isj* which have been studied in [7] and [2, 3].

These operational quantities have been applied to characterize the classes of operators in Fredholm theory: see Theorem 1 in the preliminaries. For an excellent exposition of the Fredholm theory using operational quantities we refer to Chapter 14 in Schechter's book [8].

---

*AMS Subject Classification:* 47A53

<sup>†</sup>Supported in part by DGI (Spain), Proyecto BFM2001-1147

<sup>‡</sup>Supported in part by Ministerio de Ciencia y Tecnología (Spain) and FEDER, BFM2003-07139.

From the minimum modulus  $\gamma$  we can derive eight quantities  $is\gamma$ ,  $sis\gamma$ ,  $i^*s\gamma$ ,  $s\gamma$ ,  $i\gamma$ ,  $s^*i\gamma$ ,  $isi\gamma$  and  $si\gamma$ . In this paper we obtain that  $i\gamma$  agrees with the injection modulus  $j$  and we derive, consequently, that the quantities  $s^*i\gamma$ ,  $isi\gamma$  and  $si\gamma$  are the known quantities  $s^*j$ ,  $isj$  and  $sj$ , respectively.

On the other hand, we prove that if the operator  $T$  has finite dimensional null space  $N(T)$ , then  $sj(T) \leq s\gamma(T) \leq 2sj(T)$  and we obtain that  $s\gamma$ ,  $sis\gamma$  and  $i^*s\gamma$  characterize the strictly singular operators, while  $is\gamma$  characterizes the upper semi-Fredholm operators. In the case  $N(T)$  infinite dimensional, we have  $s\gamma(T) = n(T)$  and we obtain that the quantities  $is\gamma$ ,  $sis\gamma$  and  $i^*s\gamma$  characterize the upper semi-Fredholm operators, the strictly singular operators and the compact operators, respectively.

Along the paper,  $X$ ,  $Y$ ,  $Z$  and  $W$  are infinite dimensional Banach spaces. By  $L(X, Y)$  we denote the space of all (linear continuous) operators from  $X$  into  $Y$ . For a (closed) subspace  $M$  of  $X$ ,  $J_M$  is the inclusion of  $M$  into  $X$ . An operator  $T \in L(X, Y)$  is *upper semi-Fredholm* if its range is closed and its null space is finite dimensional; it is *strictly singular* if no restriction  $TJ_M$  of  $T$  to a closed infinite dimensional subspace  $M$  of  $X$  is an isomorphism.

## 2. Preliminaries

Roughly speaking, an operational quantity is a procedure  $a$  which determines a real number  $a(T) \geq 0$  for every operator  $T$ . For two quantities  $a$  and  $b$  we write  $a \leq b$  when

$$a(T) \leq b(T), \text{ for every operator } T.$$

We say that the operational quantities  $a$  and  $b$  are *equivalent* if  $\alpha a \leq b \leq \beta a$ , for some  $\beta > \alpha > 0$ .

Given an operational quantity  $a$  and denoting by  $J_M$  the canonical inclusion of  $M$  into  $X$ , for every operator  $T \in L(X, Y)$ , where  $X$  is an infinite dimensional space, we derive the following basic quantities:

$$\begin{aligned} s^*a(T) &:= \sup\{a(TJ_P) : P \text{ finite codimensional subspace of } X\}, \\ sa(T) &:= \sup\{a(TJ_M) : M \text{ infinite dimensional subspace of } X\}, \\ i^*a(T) &:= \inf\{a(TJ_P) : P \text{ finite codimensional subspace of } X\}, \\ ia(T) &:= \inf\{a(TJ_M) : M \text{ infinite dimensional subspace of } X\}. \end{aligned}$$

Repeating the procedure, we could derive new quantities like  $sia$ ,  $sis$ ,  $i^*issa$ ,  $\dots$ , but surprisingly we obtain only three different new quantities when  $a$  is monotone [5]:

If  $a$  is *increasing*, in the sense that  $a(TJ_M) \leq a(T)$  for every  $M$ , then  $ia$ ,  $sia$  and  $i^*a$  are the only new quantities, and they satisfy

$$ia \leq sia \leq i^*a \leq a.$$

If  $a$  is *decreasing*, in the sense that  $a(TJ_M) \geq a(T)$  for every  $M$ , then  $sa$ ,  $isa$  and  $s^*a$  are the only new quantities, and they satisfy

$$a \leq s^*a \leq isa \leq sa.$$

The norm  $n(T) \equiv \|T\|$  is an increasing quantity. So we get  $i^*n$ ,  $in$  and  $sin$ . The injection modulus  $j(T)$  is decreasing. So we get  $s^*j$ ,  $sj$  and  $isj$ . There are some relations between these quantities:  $isj \leq in$  and  $sj \leq sin$ .

The operational quantities associated to  $n$  and  $j$  have been applied to characterize the classes of operators in Fredholm theory:

THEOREM 1. [2, 3], [7], [9]

1.  $i^*n(T) = 0 \Leftrightarrow T$  compact,
2.  $sin(T) = 0 \Leftrightarrow sj(T) = 0 \Leftrightarrow T$  strictly singular,
3.  $in(T) > 0 \Leftrightarrow s^*j(T) > 0 \Leftrightarrow isj(T) > 0 \Leftrightarrow T$  upper semi-Fredholm.

The quantity  $isj$  has been introduced in [2]; moreover, it was proved in [3] that, although

$$s^*j \leq isj \leq in,$$

these quantities are pairwise non-equivalent.

### 3. Main results

The operational quantity  $\gamma$  is not monotone, but  $i\gamma$  is decreasing and  $s\gamma$  is increasing. Hence we derive from  $\gamma$  the quantities

$$\begin{aligned} i\gamma &\leq s^*i\gamma \leq isi\gamma \leq si\gamma; \\ is\gamma &\leq sis\gamma \leq i^*s\gamma \leq s\gamma. \end{aligned}$$

We begin by studying the operational quantities associated with  $i\gamma$ .

THEOREM 2. For every  $T \in L(X, Y)$ ,  $i\gamma(T) = j(T)$ .

*Proof.* Let  $T$  in  $L(X, Y)$ . If  $N(T) = \{0\}$  or  $N(T)$  is infinite dimensional, then the statement is obvious. So we assume that  $0 < \dim N(T) < \infty$ , hence  $j(T) = 0$ .

We choose  $x \in N(T)$  with  $\|x\| = 1$  and  $y \in X \setminus N(T)$ , and we write

$$X = N(T) \oplus \langle y \rangle \oplus M,$$

where  $\langle y \rangle$  is the subspace generated by  $y$  and  $M$  is a closed complement of  $N(T) \oplus \langle y \rangle$ . Moreover, denoting  $y_n = (1/n)y$ , we define

$$M_n := M \oplus \langle x + y_n \rangle.$$

Suppose that  $z \in M_n \cap N(T)$  then

$$z = m + \lambda(x + y_n) = \lambda x + \lambda y_n + m,$$

for some  $m \in M$  and some scalar  $\lambda$ . Thus  $\lambda y_n = m = 0$ , and we conclude  $z = 0$ .

Since  $TJ_{M_n}$  is injective,  $\|x + y_n\| \rightarrow 1$  and  $\|T(x + y_n)\| = (1/n)\|Ty\| \rightarrow 0$ , we have  $\gamma(TJ_{M_n}) \rightarrow 0$ ; hence  $i\gamma(T) = 0$ . ■

COROLLARY 1. Let  $T \in L(X, Y)$ .

1.  $i\gamma(T) > 0 \Leftrightarrow T$  isomorphism (into);
2.  $si\gamma(T) = 0 \Leftrightarrow T$  strictly singular;
3.  $isi\gamma(T) > 0 \Leftrightarrow s^*i\gamma(T) > 0 \Leftrightarrow T$  upper semi-Fredholm.

*Proof.* From  $i\gamma = j$  we derive  $s^*i\gamma = s^*j$ ,  $isi\gamma = isj$  and  $si\gamma = sj$ . From Theorem 1 we obtain the statement. ■

Now we study the quantities associated to  $s\gamma$ . We will see that the behavior of some of these quantities depends largely on whether the null space of  $T$  is finite dimensional or infinite dimensional.

LEMMA 1. Let  $N$  be a finite dimensional subspace of  $X$ , and let  $0 < \varepsilon < 1$ . For every infinite dimensional subspace  $M$  of  $X$ , there exists a finite codimensional subspace  $M_\varepsilon$  of  $M$  such that, for every  $x \in M_\varepsilon$ ,

$$\|x\| \leq (2 + \varepsilon) \text{dist}(x, N).$$

*Proof.* Let  $\{y_1, \dots, y_k\}$  be an  $(\varepsilon/2)$ -net in the unit sphere of  $N$ . We choose  $f_1, \dots, f_k$  in the unit sphere of the dual space  $X^*$  of  $X$  so that  $f_i(y_i) = 1$  for  $i = 1, 2, \dots, k$ , and take

$$M_\varepsilon := \{x \in M : f_1(x) = \dots = f_k(x) = 0\}.$$

Let  $x \in M_\varepsilon$ . For each  $y \in N$  we denote  $z_i := \|y\|y_i$  ( $1 \leq i \leq k$ ). Then

$$\|y - x\| \geq \|x - z_i\| - \|y - z_i\| \geq f_i(z_i) - (\varepsilon/2)\|y\| = \|y\| - (\varepsilon/2)\|y\| \geq \frac{\|y\|}{1 + \varepsilon}$$

for some  $i$ . Hence,  $\|y\| \leq (1 + \varepsilon)\|y - x\|$  for each  $y$  in  $N$ . From this we obtain

$$\|x\| \leq \|x - y\| + \|y\| \leq (2 + \varepsilon)\|x - y\|,$$

for each  $y$  in  $N$ , and this implies  $\|x\| \leq (2 + \varepsilon) \text{dist}(x, N)$ . ■

THEOREM 3. Let  $T \in L(X, Y)$ . If  $N(T)$  is finite dimensional, then

$$sj(T) \leq s\gamma(T) \leq 2sj(T).$$

*Proof.* Let  $M$  be an infinite dimensional subspace of  $X$ , and let  $\varepsilon > 0$ . By Lemma 1, there exists a finite codimensional subspace  $M_\varepsilon$  of  $M$  such that for  $x \in M_\varepsilon$ ,

$$\|x\| \leq (2 + \varepsilon) \text{dist}(x, N(T)) \leq (2 + \varepsilon) \text{dist}(x, N(TJ_M)).$$

Then

$$\begin{aligned} \gamma(TJ_M) &= \inf_{x \in M, Tx \neq 0} \frac{\|Tx\|}{\text{dist}(x, N(TJ_M))} \\ &\leq \inf_{x \in M_\varepsilon, Tx \neq 0} \frac{\|Tx\|}{\text{dist}(x, N(TJ_M))} \\ &\leq \inf_{x \in M_\varepsilon, Tx \neq 0} \frac{\|Tx\|}{\|x\|} (2 + \varepsilon) = (2 + \varepsilon)j(TJ_{M_\varepsilon}). \end{aligned}$$

Hence,  $\gamma(TJ_M) \leq (2 + \varepsilon)sj(T)$ , so  $s\gamma(T) \leq 2sj(T)$ .

The inequality  $sj(T) \leq s\gamma(T)$  is obvious. ■

**COROLLARY 2.** *Let  $T \in L(X, Y)$ . If  $N(T)$  is finite dimensional, then*

1.  $isj(T) \leq is\gamma(T) \leq 2isj(T)$ ,
2.  $sj(T) \leq sis\gamma(T) \leq 2sj(T)$ ,
3.  $sj(T) \leq i^*s\gamma(T) \leq 2sj(T)$ .

*Proof.* Note that  $N(TJ_M)$  is also finite dimensional for each  $M$ .

1. For every infinite dimensional subspace  $M$  of  $X$  we have that  $sj(TJ_M) \leq s\gamma(TJ_M) \leq 2sj(TJ_M)$ , hence  $isj(T) \leq is\gamma(T) \leq 2isj(T)$ .

2. Similar to 1., taking into account  $sisj(T) = sj(T)$  [5].

3. Similar to 1., taking into account  $i^*sj(T) = sj(T)$  [5]. ■

The following result shows that the property  $is\gamma(T) > 0$  is preserved by taking products.

**COROLLARY 3.** *Let  $T \in L(X, Y)$  and  $S \in L(Y, Z)$ . If  $N(T)$  and  $N(S)$  are finite dimensional, then*

$$is\gamma(S) is\gamma(T) \leq 4 is\gamma(ST)$$

*Proof.* It is known that  $isj(S) isj(T) \leq isj(ST)$  [2], hence  $is\gamma(S) is\gamma(T) \leq 2isj(S) 2isj(T) \leq 4isj(ST) \leq 4is\gamma(ST)$ . ■

**THEOREM 4.** *For each operator  $T \in L(X, Y)$  with  $N(T)$  infinite dimensional,  $s\gamma(T) = n(T)$ .*

*Proof.* For  $T = 0$  the result is obvious. Suppose  $T \neq 0$ . For each  $x \notin N(T)$  we put  $M_x := N(T) \oplus \langle x \rangle$ , where  $\langle x \rangle$  is the subspace generated by  $x$ .

For every  $y \in M_x$ ,  $y = \lambda x + z$ ,  $z \in N(T)$ , we obtain

$$\frac{\|Tx\|}{\|x\|} = \frac{|\lambda| \|Tx\|}{|\lambda| \|x\|} = \frac{\|Ty\|}{\|y - z\|} \leq \frac{\|Ty\|}{\text{dist}(y, N(T))}.$$

Thus  $n(T) \leq \gamma(TJ_{M_x}) \leq s\gamma(T)$ . ■

Note that for every  $T \neq 0$  with  $R(T)$  finite dimensional (hence  $N(T)$  is infinite dimensional),  $sj(T) = 0 \neq n(T)$ . Thus  $sj$  and  $s\gamma$  are not equivalent.

**COROLLARY 4.** *Let  $T \in L(X, Y)$ .*

1.  $is\gamma(T) > 0 \Leftrightarrow T$  is upper semi-Fredholm.
2. If  $N(T)$  is finite dimensional, then

$$s\gamma(T) = 0 \Leftrightarrow sis\gamma(T) = 0 \Leftrightarrow i^*s\gamma(T) = 0 \Leftrightarrow T \text{ is strictly singular.}$$

3. If  $N(T)$  is infinite dimensional, then

- (a)  $s\gamma(T) = 0 \Leftrightarrow T = 0$ ;  
 (b)  $sis\gamma(T) = 0 \Leftrightarrow T$  is strictly singular;  
 (c)  $i^*s\gamma(T) = 0 \Leftrightarrow T$  is compact.

*Proof.* (1) [6, Example 5.1].

(2) It is immediate from Theorem 1, Theorem 3 and Corollary 2.

(3) (a) Theorem 4.

(b)  $sis\gamma(T) = 0$  is equivalent to  $is\gamma(TJ_M) = 0$  for every infinite dimensional subspace  $M$  of  $X$ , which is equivalent by (1) to  $TJ_M$  is not an upper semi-Fredholm operator, and consequently to  $T$  strictly singular.

(c) Since  $i^*s\gamma(T)$  is the infimum of  $s\gamma(TJ_P) = n(TJ_P)$  where  $P$  runs over the finite codimensional subspaces of  $X$ , from Theorem 1, we obtain  $i^*s\gamma(T) = i^*n(T) = 0$  if and only if  $T$  is compact. ■

#### REFERENCES

- [1] S. Goldberg, *Unbounded Linear Operators*, McGraw-Hill, New York, 1966 [Dover, New York, 1985].  
 [2] M. González, A. Martínón, *Fredholm Theory and Space Ideals*, Bol. U.M.I. (7) 7-B (1993), 473–488.  
 [3] M. González, A. Martínón, *Operational quantities characterizing semi-Fredholm operators*, Studia Math. 114 (1995), 13–27.  
 [4] A. Lebow, M. Schechter, *Semigroups of operators and measures of noncompactness*, J. Funct. Anal. 7 (1971), 1–26.  
 [5] A. Martínón, *Generating real maps on a biordered set*, Comment. Math. Univ. Carolinae 32 (1991), 265–272.  
 [6] A. Martínón, *Operational quantities*, Comment. Math. Univ. Carolinae 38 (1997), 471–484.  
 [7] M. Schechter, *Quantities related to strictly singular operators*, Indiana Univ. Math. J. 21 (1972), 1061–1071.  
 [8] M. Schechter, *Principles of Functional Analysis*, American Mathematical Society, Providence, Rhode Island, 2002.  
 [9] A. A. Sedaev, *The structure of certain linear operators* (in Russian), Mat. Issled. 5 (1970), 166–175.

(received 11.02.2003, in revised form 23.12.2004)

Manuel González, Departamento de Matemáticas, Universidad de Cantabria, 39071 Santander, Spain

*E-mail:* gonzalem@unican.es

Antonio Martínón, Departamento de Análisis Matemático, Universidad de La Laguna, 38271 La Laguna (Tenerife), Spain

*E-mail:* anmarce@ull.es