

Full faithfulness theorem for torsion crystalline representations

Yoshiyasu Ozeki

ABSTRACT. Mark Kisin proved that a certain “restriction functor” on crystalline p -adic representations is fully faithful. In this paper, we prove the torsion analogue of Kisin’s theorem.

CONTENTS

1. Introduction	1043
2. Preliminaries	1044
3. Proof of Theorem 1	1047
4. Nonfullness: Examples	1053
Appendix A. (φ, \hat{G}) -modules associated with crystalline representations	1057
References	1060

1. Introduction

Let $p > 2$ be a prime number and $r, r' \geq 0$ integers. Let K be a complete discrete valuation field of mixed characteristic $(0, p)$ with perfect residue field and absolute ramification index e . Let $\pi = \pi_0$ be a uniformizer of K and π_n a p^n -th root of π such that $\pi_{n+1}^p = \pi_n$ for all $n \geq 0$. Put $K_\infty = \bigcup_{n \geq 0} K(\pi_n)$ and denote by G_K and G_∞ absolute Galois groups of K and K_∞ , respectively. In Theorem (0.2) of [Kis], Kisin proved that the functor “restriction to G_∞ ” from crystalline \mathbb{Q}_p -representations of G_K to \mathbb{Q}_p -representations of G_∞ is fully faithful, which was a conjecture of Breuil ([Br2]). Hence we may say that crystalline \mathbb{Q}_p -representations of G_K are characterized by their restriction to G_∞ . It should be noted that there exists an established theory describing representations of G_∞ by easy linear algebra data, which is called étale φ -modules, introduced by Fontaine ([Fo1] A 1.2). In this paper, we are interested in the torsion analogue of the above Kisin’s result. For example, Breuil proved in Theorem 3.4.3 of [Br3] that

Received October 7, 2014.

2010 *Mathematics Subject Classification.* 11F85, 11S20.

Key words and phrases. Torsion crystalline representations, fully faithful.

This work was supported by JSPS KAKENHI Grant Number 25-173.

the functor “restriction to G_∞ ” from finite flat representations of G_K to torsion \mathbb{Z}_p -representations of G_∞ is fully faithful (Remark 20(2)). Our main theorem is motivated by his result:

Theorem 1. *Suppose $er < p - 1$ and $e(r' - 1) < p - 1$. Let T (resp. T') be a torsion crystalline \mathbb{Z}_p -representation of G_K with Hodge–Tate weights in $[0, r]$ (resp. $[0, r']$). Then any G_∞ -equivariant morphism $T \rightarrow T'$ is in fact G_K -equivariant.*

In particular, the functor from torsion crystalline \mathbb{Z}_p -representations of G_K with Hodge–Tate weights in $[0, r]$ to torsion \mathbb{Z}_p -representations of G_∞ , obtained by restricting the action of G_K to G_∞ , is fully faithful.

Here a torsion \mathbb{Z}_p -representation of G_K is said to be *torsion crystalline with Hodge–Tate weights in $[0, r]$* if it can be written as the quotient of two lattices in some crystalline \mathbb{Q}_p -representation of G_K with Hodge–Tate weights in $[0, r]$. For example, a torsion \mathbb{Z}_p -representation of G_K is finite flat if and only if it is torsion crystalline with Hodge–Tate weights in $[0, 1]$ (Remark 20(2)). If $e = 1$, the latter part of Theorem 1 has been proven by Breuil via Fontaine–Laffaille theory (Remark 20(3)). On the other hand, our proof is based on results on Kisin modules and (φ, \hat{G}) -modules (the notion of (φ, \hat{G}) -modules is introduced in [Li2]). More precisely, we use maximal models for Kisin modules introduced in [CL1] and results on “the range of monodromy” for (φ, \hat{G}) -modules given in Section 4 of [GLS].

It seems natural to ask whether the condition “ $er < p - 1$ ” in the latter part of Theorem 1 is necessary and sufficient for the full faithfulness. In fact, we know that the condition “ $er < p - 1$ ” is not necessary since our restriction functor is fully faithful for any e when $r = 1$ (Remark 20(2)). (Maybe the necessary and sufficient condition for the full faithfulness is “ $e(r - 1) < p - 1$ ” (Remark 20).) In addition, in the last section, we give some examples such that the restriction functor appeared in Theorem 1 is not full under some choices of K and r which do not satisfy “ $er < p - 1$ ” (more precisely, “ $e(r - 1) < p - 1$ ”). Examples are mainly given by using two methods: The first one is direct computations of Galois cohomologies, which is a purely local method. The second one is based on the classical Serre’s modularity conjecture, which is a global method.

Acknowledgements. It is a pleasure to thank Wansu Kim for useful comments and correspondences to Theorem 1. The author thanks Naoki Imai and Akio Tamagawa who gave him useful advice in the proof of his main theorem. The author thanks also Keisuke Arai, Seidai Yasuda, Shin Hattori and Yuichiro Taguchi for their helpful comments on Proposition 16.

2. Preliminaries

Throughout this paper, we fix a prime number $p > 2$. Let $r \geq 0$ be an integer. Let k be a perfect field of characteristic p , $W(k)$ its ring of Witt

vectors, $K_0 = W(k)[1/p]$, K a finite totally ramified extension of K_0 , \bar{K} a fixed algebraic closure of K and $G_K = \text{Gal}(\bar{K}/K)$. Fix a uniformizer $\pi \in K$ and denote by $E(u)$ its Eisenstein polynomial over K_0 . For any integer $n \geq 0$, let $\pi_n \in \bar{K}$ be a p^n -th root of π such that $\pi_{n+1}^p = \pi_n$. Let $K_\infty = \bigcup_{n \geq 0} K(\pi_n)$ and $G_\infty = \text{Gal}(\bar{K}/K_\infty)$.

For any topological group H , we denote by $\text{Rep}_{\text{tor}}(H)$ (resp. $\text{Rep}_{\mathbb{Z}_p}(H)$) the category of finite torsion \mathbb{Z}_p -representations of H (resp. the category of finite free \mathbb{Z}_p -representations of H). We denote by $\text{Rep}_{\mathbb{Z}_p}^r(G_K)$ the category of lattices in crystalline \mathbb{Q}_p -representations of G_K with Hodge–Tate weights in $[0, r]$. We say that $T \in \text{Rep}_{\text{tor}}(G_K)$ is *torsion crystalline with Hodge–Tate weights in $[0, r]$* if it can be written as the quotient of $L' \subset L$ in $\text{Rep}_{\mathbb{Z}_p}^r(G_K)$, and denote by $\text{Rep}_{\text{tor}}^r(G_K)$ the category of them.

Let $R = \varprojlim \mathcal{O}_{\bar{K}}/p$ where $\mathcal{O}_{\bar{K}}$ is the integer ring of \bar{K} and the transition maps are given by the p -th power map. Write $\underline{\pi} = (\pi_n)_{n \geq 0} \in R$ and let $[\underline{\pi}] \in W(R)$ be the Teichmüller representative of $\underline{\pi}$. Let $\mathfrak{S} = W(k)[[u]]$ equipped with a Frobenius endomorphism φ given by $u \mapsto u^p$ and the Frobenius on $W(k)$. We embed the $W(k)$ -algebra $W(k)[u]$ into $W(R)$ via the map $u \mapsto [\underline{\pi}]$. This embedding extends to an embedding $\mathfrak{S} \hookrightarrow W(R)$, which is compatible with Frobenius endomorphisms.

A φ -module (over \mathfrak{S}) is an \mathfrak{S} -module \mathfrak{M} equipped with a φ -semilinear map $\varphi: \mathfrak{M} \rightarrow \mathfrak{M}$. A morphism between two φ -modules $(\mathfrak{M}_1, \varphi_1)$ and $(\mathfrak{M}_2, \varphi_2)$ is an \mathfrak{S} -linear map $\mathfrak{M}_1 \rightarrow \mathfrak{M}_2$ compatible with φ_1 and φ_2 . Denote by $'\text{Mod}_{/\mathfrak{S}}^r$ the category of φ -modules (\mathfrak{M}, φ) of height $\leq r$ in the sense that \mathfrak{M} is of finite type over \mathfrak{S} and the cokernel of $1 \otimes \varphi: \mathfrak{S} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M} \rightarrow \mathfrak{M}$ is killed by $E(u)^r$. Let $\text{Mod}_{/\mathfrak{S}_\infty}^r$ be the full subcategory of $'\text{Mod}_{/\mathfrak{S}}^r$ consisting of finite \mathfrak{S} -modules which are killed by some power of p and have projective dimension 1 in the sense that \mathfrak{M} has a two term resolution by finite free \mathfrak{S} -modules. Let $\text{Mod}_{/\mathfrak{S}}^r$ be the full subcategory of $'\text{Mod}_{/\mathfrak{S}}^r$ consisting of finite free \mathfrak{S} -modules. We call an object of $\text{Mod}_{/\mathfrak{S}_\infty}^r$ (resp. $\text{Mod}_{/\mathfrak{S}}^r$) a *torsion Kisin module* (resp. a *free Kisin module*). A *Kisin module* is a torsion Kisin module or a free Kisin module. For any Kisin module \mathfrak{M} , we define a \mathbb{Z}_p -representation $T_{\mathfrak{S}}(\mathfrak{M})$ of G_∞ by

$$T_{\mathfrak{S}}(\mathfrak{M}) = \begin{cases} \text{Hom}_{\mathfrak{S}, \varphi}(\mathfrak{M}, \mathbb{Q}_p/\mathbb{Z}_p \otimes_{\mathbb{Z}_p} \mathfrak{S}^{\text{ur}}) & \text{if } \mathfrak{M} \text{ is torsion} \\ \text{Hom}_{\mathfrak{S}, \varphi}(\mathfrak{M}, \mathfrak{S}^{\text{ur}}) & \text{if } \mathfrak{M} \text{ is free.} \end{cases}$$

Here, a G_∞ -action on $T_{\mathfrak{S}}(\mathfrak{M})$ is given by $(\sigma.f)(x) = \sigma(f(x))$ for $\sigma \in G_\infty, f \in T_{\mathfrak{S}}(\mathfrak{M}), x \in \mathfrak{M}$.

Here we recall the theory of Liu’s (φ, \hat{G}) -modules (cf. [Li2]). Let S be the p -adic completion of the divided power envelope of $W(k)[u]$ with respect to the ideal generated by $E(u)$. There exists a unique Frobenius map $\varphi: S \rightarrow S$ defined by $\varphi(u) = u^p$. Put $S_{K_0} = S[1/p] = K_0 \otimes_{W(k)} S$. The inclusion $W(k)[u] \hookrightarrow W(R)$ via the map $u \mapsto [\underline{\pi}]$ induces φ -compatible inclusions $\mathfrak{S} \hookrightarrow S \hookrightarrow A_{\text{cris}}$ and $S_{K_0} \hookrightarrow B_{\text{cris}}^+$. Fix a choice of primitive p^i -root of

unity ζ_{p^i} for $i \geq 0$ such that $\zeta_{p^{i+1}}^p = \zeta_{p^i}$. Put $\varepsilon = (\zeta_{p^i})_{i \geq 0} \in R^\times$ and $t = \log([\varepsilon]) \in A_{\text{cris}}$. Denote by $\nu: W(R) \rightarrow W(\bar{k})$ a unique lift of the projection $R \rightarrow \bar{k}$, which extends to a map $\nu: B_{\text{cris}}^+ \rightarrow W(\bar{k})[1/p]$. For any subring $A \subset B_{\text{cris}}^+$, we put $I_+A = \text{Ker}(\nu \text{ on } B_{\text{cris}}^+) \cap A$. For any integer $n \geq 0$, let $t^{\{n\}} = t^{r(n)} \gamma_{\tilde{q}(n)}(\frac{t^{p-1}}{p})$ where $n = (p-1)\tilde{q}(n) + r(n)$ with $\tilde{q}(n) \geq 0$, $0 \leq r(n) < p-1$ and $\gamma_i(x) = \frac{x^i}{i!}$ is the standard divided power. We define a subring \mathcal{R}_{K_0} of B_{cris}^+ as below:

$$\mathcal{R}_{K_0} = \left\{ \sum_{i=0}^{\infty} f_i t^{\{i\}} \mid f_i \in S_{K_0} \text{ and } f_i \rightarrow 0 \text{ as } i \rightarrow \infty \right\}.$$

Put $\widehat{\mathcal{R}} = \mathcal{R}_{K_0} \cap W(R)$ and $I_+ = I_+ \widehat{\mathcal{R}}$. Put $\widehat{K} = \bigcup_{n \geq 0} K_\infty(\zeta_{p^n})$ and $\widehat{G} = \text{Gal}(\widehat{K}/K)$. Lemma 2.2.1 in [Li2] shows that $\widehat{\mathcal{R}}$ (resp. \mathcal{R}_{K_0}) is a φ -stable \mathfrak{S} -algebra as a subring in $W(R)$ (resp. B_{cris}^+), and ν induces $\mathcal{R}_{K_0}/I_+ \mathcal{R}_{K_0} \simeq K_0$ and $\widehat{\mathcal{R}}/I_+ \simeq S/I_+ S \simeq \mathfrak{S}/I_+ \mathfrak{S} \simeq W(\bar{k})$. Furthermore, $\widehat{\mathcal{R}}, I_+, \mathcal{R}_{K_0}$ and $I_+ \mathcal{R}_{K_0}$ are G_K -stable, and G_K -actions on them factors through \widehat{G} . For any Kisin module \mathfrak{M} , we equip $\widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$ with a Frobenius by $\varphi_{\widehat{\mathcal{R}}} \otimes \varphi_{\mathfrak{M}}$. It is known that the natural map $\mathfrak{M} \rightarrow \widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$ given by $x \mapsto 1 \otimes x$ is an injection ([CL2], Section 3.1). By this injection, we regard \mathfrak{M} as a $\varphi(\mathfrak{S})$ -stable submodule of $\widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$.

Definition 2. A (φ, \widehat{G}) -module (of height $\leq r$) is a triple $\widehat{\mathfrak{M}} = (\mathfrak{M}, \varphi_{\mathfrak{M}}, \widehat{G})$ where:

- (1) $(\mathfrak{M}, \varphi_{\mathfrak{M}})$ is a Kisin module (of height $\leq r$).
- (2) \widehat{G} is a continuous $\widehat{\mathcal{R}}$ -semilinear \widehat{G} -action on $\widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$.
- (3) the \widehat{G} -action commutes with $\varphi_{\widehat{\mathcal{R}}} \otimes \varphi_{\mathfrak{M}}$.
- (4) $\mathfrak{M} \subset (\widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M})^{H_K}$ where $H_K = \text{Gal}(\widehat{K}/K_\infty)$.
- (5) \widehat{G} acts on the $W(\bar{k})$ -module $(\widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M})/I_+(\widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M})$ trivially.

If \mathfrak{M} is a torsion (resp. free) Kisin module, we call $\widehat{\mathfrak{M}}$ a *torsion* (resp. *free*) (φ, \widehat{G}) -module.

A morphism between two (φ, \widehat{G}) -modules $\widehat{\mathfrak{M}}_1 = (\mathfrak{M}_1, \varphi_1, \widehat{G})$ and $\widehat{\mathfrak{M}}_2 = (\mathfrak{M}_2, \varphi_2, \widehat{G})$ is a morphism $f: \mathfrak{M}_1 \rightarrow \mathfrak{M}_2$ of φ -modules such that

$$\widehat{\mathcal{R}} \otimes f: \widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}_1 \rightarrow \widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}_2$$

is \widehat{G} -equivariant. We denote by $\text{Mod}_{\mathfrak{S}_\infty}^{r, \widehat{G}}$ (resp. $\text{Mod}_{\mathfrak{S}}^{r, \widehat{G}}$) the category of torsion (φ, \widehat{G}) -modules of height $\leq r$ (resp. free (φ, \widehat{G}) -modules of height $\leq r$). We often regard $\widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$ as a G_K -module via the projection $G_K \rightarrow \widehat{G}$. A sequence $0 \rightarrow \widehat{\mathfrak{M}}' \rightarrow \widehat{\mathfrak{M}} \rightarrow \widehat{\mathfrak{M}}'' \rightarrow 0$ of (φ, \widehat{G}) -modules is *exact* if it is exact as \mathfrak{S} -modules. For a (φ, \widehat{G}) -module $\widehat{\mathfrak{M}}$, we define a \mathbb{Z}_p -representation

$\hat{T}(\hat{\mathfrak{M}})$ of G_K by

$$\hat{T}(\hat{\mathfrak{M}}) = \begin{cases} \text{Hom}_{\hat{\mathcal{R}},\varphi}(\hat{\mathcal{R}} \otimes_{\varphi,\mathfrak{S}} \mathfrak{M}, \mathbb{Q}_p/\mathbb{Z}_p \otimes_{\mathbb{Z}_p} W(R)) & \text{if } \mathfrak{M} \text{ is torsion} \\ \text{Hom}_{\hat{\mathcal{R}},\varphi}(\hat{\mathcal{R}} \otimes_{\varphi,\mathfrak{S}} \mathfrak{M}, W(R)) & \text{if } \mathfrak{M} \text{ is free.} \end{cases}$$

Here, G_K acts on $\hat{T}(\hat{\mathfrak{M}})$ by $(\sigma.f)(x) = \sigma(f(\sigma^{-1}(x)))$ for $\sigma \in G_K$, $f \in \hat{T}(\hat{\mathfrak{M}})$, $x \in \hat{\mathcal{R}} \otimes_{\varphi,\mathfrak{S}} \mathfrak{M}$. Then, there exists a natural G_∞ -equivariant map

$$\theta: T_{\mathfrak{S}}(\mathfrak{M}) \rightarrow \hat{T}(\hat{\mathfrak{M}})$$

defined by $\theta(f)(a \otimes m) = a\varphi(f(m))$ for $f \in T_{\mathfrak{S}}(\mathfrak{M})$, $a \in \hat{\mathcal{R}}$, $m \in \mathfrak{M}$.

Fix a topological generator τ of $\text{Gal}(\hat{K}/K_{p^\infty})$ where $K_{p^\infty} = \bigcup_{n \geq 0} K(\zeta_{p^n})$. We may suppose that $\zeta_{p^n} = \tau(\pi_n)/\pi_n$ for all n , and this implies $\tau(u) = [\varepsilon]u$ in $W(R)$. There exists $\mathfrak{t} \in W(R) \setminus pW(R)$ such that $\varphi(\mathfrak{t}) = pE(0)^{-1}E(u)\mathfrak{t}$. Such \mathfrak{t} is unique up to units of \mathbb{Z}_p (cf. Example 2.3.5 of [Li1]). The following theorems play important rolls in the proof of Theorem 1.

Theorem 3 ([Li2]).

- (1) *The map $\theta: T_{\mathfrak{S}}(\mathfrak{M}) \rightarrow \hat{T}(\hat{\mathfrak{M}})$ is an isomorphism.*
- (2) *The contravariant functor \hat{T} induces an anti-equivalence between the category $\text{Mod}_{\mathfrak{S}}^{r,\hat{G}}$ of free (φ, \hat{G}) -modules of height $\leq r$ and the category of G_K -stable \mathbb{Z}_p -lattices in semi-stable \mathbb{Q}_p -representations of G_K with Hodge–Tate weights in $[0, r]$.*

Theorem 4 ([CL2], Theorem 3.1.3 (4), [GLS], Proposition 5.9). *Let $T \in \text{Rep}_{\text{tor}}^r(G_K)$ and take $L' \subset L$ in $\text{Rep}_{\mathbb{Z}_p}^r(G_K)$ such that $T \simeq L/L'$.*

- (1) *There exists an exact sequence $\mathfrak{S}: 0 \rightarrow \hat{\mathfrak{L}} \rightarrow \hat{\mathfrak{L}}' \rightarrow \hat{\mathfrak{M}} \rightarrow 0$ of (φ, \hat{G}) -modules such that:*
 - (a) *$\hat{\mathfrak{L}}$ and $\hat{\mathfrak{L}}'$ are free (φ, \hat{G}) -modules of height $\leq r$.*
 - (b) *$\hat{\mathfrak{M}}$ is a torsion (φ, \hat{G}) -module of height $\leq r$.*
 - (c) *$\hat{T}(\mathfrak{S})$ is isomorphic to the exact sequence $0 \rightarrow L' \rightarrow L \rightarrow T \rightarrow 0$ of $\mathbb{Z}_p[G_K]$ -modules.*
- (2) *Let $\hat{\mathfrak{M}}$ be as in (1). For any $x \in \mathfrak{M}$, we have*

$$\tau(x) - x \in u^p \varphi(\mathfrak{t})(W(R) \otimes_{\varphi,\mathfrak{S}} \mathfrak{M}).$$

Proof. The assertion (2) is an easy consequence of [GLS], Proposition 5.9. Here is one remark: In *loc. cit.*, K is assumed to be a finite extension of \mathbb{Q}_p , but arguments in Section 4.1 and 4.2 of *loc. cit.* proceed even if K is not only a finite extension of \mathbb{Q}_p but also any complete discrete valuation field of mixed characteristic $(0, p)$ with perfect residue field. \square

3. Proof of Theorem 1

For any integer $\alpha \geq 0$, we denote by $\mathfrak{m}_R^{\geq \alpha}$ the ideal of R consisting of $a \in R$ with $v_R(a) \geq \alpha$, where v_R is a valuation of R such that $v_R(\pi) = \frac{1}{e}$.

Note that, if we put $\tilde{\mathfrak{t}} = \mathfrak{t} \bmod p \in R$, then $v_R(\tilde{\mathfrak{t}}) = \frac{1}{p-1}$ since $\varphi(\tilde{\mathfrak{t}}) \in \pi^e \tilde{\mathfrak{t}} \cdot R^\times$ (recall the equation $\varphi(\mathfrak{t}) = pE(0)^{-1}E(u)\mathfrak{t}$).

We note that we have natural inclusions

$$\mathfrak{M} \subset \mathfrak{S} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M} \subset \hat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M} \subset W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$$

for any $\mathfrak{M} \in \text{Mod}_{/\mathfrak{S}_\infty}^r$. Denote by $\text{Mod}_{/\mathfrak{S}_\infty}^{r, \hat{G}, \text{cris}}$ the full subcategory of $\text{Mod}_{/\mathfrak{S}_\infty}^{r, \hat{G}}$ consisting of torsion (φ, \hat{G}) -modules \mathfrak{M} which satisfy the following; for any $x \in \mathfrak{M}$,

$$\tau(x) - x \in u^p \varphi(\mathfrak{t})(W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}).$$

We define the full subcategory $\text{Rep}_{\text{tor}}^{r, \hat{G}, \text{cris}}(G_K)$ of $\text{Rep}_{\text{tor}}(G_K)$ to be the essential image of the functor $\text{Mod}_{/\mathfrak{S}_\infty}^{r, \hat{G}, \text{cris}} \subset \text{Mod}_{/\mathfrak{S}_\infty}^{r, \hat{G}} \xrightarrow{\hat{T}} \text{Rep}_{\text{tor}}(G_K)$, where \hat{T} is defined in the previous section. By Theorem 4, we have

$$\text{Rep}_{\text{tor}}^r(G_K) \subset \text{Rep}_{\text{tor}}^{r, \hat{G}, \text{cris}}(G_K).$$

Remark 5.

- (1) The subscript “cris” of $\text{Mod}_{/\mathfrak{S}_\infty}^{r, \hat{G}, \text{cris}}$ is plausible since a free (φ, \hat{G}) -module \mathfrak{M} satisfying the condition

$$\tau(x) - x \in u^p \varphi(\mathfrak{t})(W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M})$$

corresponds to a crystalline representation. See Theorem 21 in the appendix for more precise information.

- (2) Note that objects of $\text{Rep}_{\text{tor}}^{r, \hat{G}, \text{cris}}(G_K)$ are not necessarily torsion crystalline representations. In fact, we do not know whether torsion (φ, \hat{G}) -modules lift to free (φ, \hat{G}) -modules.

Theorem 1 follows from the following result.

Theorem 6. *Let $T \in \text{Rep}_{\text{tor}}^{r, \hat{G}, \text{cris}}(G_K)$ and $T' \in \text{Rep}_{\text{tor}}^{r', \hat{G}, \text{cris}}(G_K)$, and suppose $er < p - 1$ and $e(r' - 1) < p - 1$. Then any G_∞ -equivariant morphism $T \rightarrow T'$ is in fact G_K -equivariant.*

Lemma 7. *Let $a \in W(R) \setminus pW(R)$. For any Kisin module \mathfrak{M} , the map*

$$W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M} \rightarrow W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}, \quad x \mapsto ax$$

is injective.

Proof. We may suppose that \mathfrak{M} is a torsion Kisin module. By a dévissage argument ([Lil], Proposition 2.3.2 (4)), we may assume $p\mathfrak{M} = 0$. In this situation, the statement is clear since $W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$ is a finite direct sum of R . □

The following is a key lemma for our proof of Theorem 6:

Lemma 8. *Let r and r' be nonnegative integers with $e(r - 1) < p - 1$ (without any assumption on r'). Let $\hat{\mathfrak{M}}$ and $\hat{\mathfrak{N}}$ be objects of $\text{Mod}_{\mathfrak{S}_\infty}^{r, \hat{G}, \text{cris}}$ and $\text{Mod}_{\mathfrak{S}_\infty}^{r', \hat{G}, \text{cris}}$, respectively. Then we have $\text{Hom}(\hat{\mathfrak{M}}, \hat{\mathfrak{N}}) = \text{Hom}(\mathfrak{M}, \mathfrak{N})$.*

In particular, if $e(r - 1) < p - 1$, then the forgetful functor

$$\text{Mod}_{\mathfrak{S}_\infty}^{r, \hat{G}, \text{cris}} \rightarrow \text{Mod}_{\mathfrak{S}_\infty}^r$$

is fully faithful.

The condition $e(r - 1) < p - 1$ is essential. See Remark 11 below.

Proof. Let $f: \mathfrak{M} \rightarrow \mathfrak{N}$ be a morphism of Kisin modules and put

$$\hat{f} = W(R) \otimes f: W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M} \rightarrow W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}.$$

It suffices to prove that, for any $x \in \mathfrak{M}$, $\Delta(1 \otimes x) = 0$ where $\Delta = \tau \circ \hat{f} - \hat{f} \circ \tau$. We proceed by induction on n such that $p^n \mathfrak{N} = 0$.

Suppose $n = 1$, that is, $p\mathfrak{N} = 0$. We may identify $W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}$ with $R \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}$. Since $\Delta(1 \otimes x) = (\tau - 1)(1 \otimes f(x)) - \hat{f}((\tau - 1)(1 \otimes x))$, we obtain the following implication

$$(0) \quad \text{For any } x \in \mathfrak{M}, \quad \Delta(1 \otimes x) \in \mathfrak{m}_R^{\geq c(0)}(R \otimes_{\varphi, \mathfrak{S}} \mathfrak{N})$$

where $c(0) = \frac{p}{p-1} + \frac{p}{e}$. Note that

$$\Delta(1 \otimes E(u)^r x) = \tau(\varphi(E(u)))^r \Delta(1 \otimes x) = (\varepsilon u)^{per} \Delta(1 \otimes x) \in R \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}.$$

On the other hand, since \mathfrak{M} is of height $\leq r$, we can write $E(u)^r x = \sum_{i \geq 0} a_i \varphi(y_i)$ for some $a_i \in \mathfrak{S}$ and $y_i \in \mathfrak{M}$. Then we obtain

$$\Delta(1 \otimes E(u)^r x) = \sum_{i \geq 0} \tau(\varphi(a_i)) \varphi(\Delta(1 \otimes y_i))$$

and it is contained in $\mathfrak{m}_R^{\geq pc(0)}(R \otimes_{\varphi, \mathfrak{S}} \mathfrak{N})$ by the implication (0). Since $R \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}$ is free as an R -module, we obtain the implication

$$(1) \quad \text{For any } x \in \mathfrak{M}, \quad \Delta(1 \otimes x) \in \mathfrak{m}_R^{\geq c(1)}(R \otimes_{\varphi, \mathfrak{S}} \mathfrak{N})$$

where $c(1) = pc(0) - pr = \frac{p^2}{p-1} + \frac{p^2}{e} - pr$. By repeating the same argument, for any $s \geq 0$, we see the following implication

$$(s) \quad \text{For any } x \in \mathfrak{M}, \quad \Delta(1 \otimes x) \in \mathfrak{m}_R^{\geq c(s)}(R \otimes_{\varphi, \mathfrak{S}} \mathfrak{N})$$

where $c(s) = pc(s - 1) - pr = \frac{p^{s+1}}{p-1} + \frac{p^{s+1}}{e} - p^s r - \dots - pr$. Since $e(r - 1) < p - 1$, we know that $\mathfrak{m}_R^{\geq c(s)}$ goes to zero when $s \rightarrow \infty$ and then we obtain $\Delta(1 \otimes x) = 0$.

Suppose $n > 1$. Consider the exact sequence

$$(*) \quad 0 \rightarrow \text{Ker}(p) \rightarrow \mathfrak{N} \xrightarrow{p} p\mathfrak{N} \rightarrow 0$$

of φ -modules. By Lemma 2.3.1 and Proposition 2.3.2 of [Li1], we know that $\mathfrak{N}' := \text{Ker}(p)$ and $\mathfrak{N}'' := p\mathfrak{N}$ are in $\text{Mod}_{\mathfrak{S}_\infty}^{r'}$. Equipping $\hat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}''$ with

\hat{G} -action via the natural identification $p(\widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}) = \widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}'$, we see that \mathfrak{N}' has a structure as a (φ, \hat{G}) -module. We can also equip $\widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}'$ with \hat{G} -action via the exact sequence

$$0 \rightarrow \widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}' \rightarrow \widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{N} \rightarrow \widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}'' \rightarrow 0$$

(for the exactness, see [CL2], Lemma 3.1.2). Since the sequence

$$0 \rightarrow \widehat{\mathcal{R}}/I_+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}' \rightarrow \widehat{\mathcal{R}}/I_+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{N} \rightarrow \widehat{\mathcal{R}}/I_+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}'' \rightarrow 0$$

is also exact ([Oz], Corollary 2.11), we know that \mathfrak{N}' also has a structure as a (φ, \hat{G}) -module. Summary, we obtained an exact sequence

$$0 \rightarrow \hat{\mathfrak{N}}' \rightarrow \hat{\mathfrak{N}} \xrightarrow{p} \hat{\mathfrak{N}}'' \rightarrow 0$$

in $\text{Mod}_{\mathfrak{S}_\infty}^{r', \hat{G}}$ whose underlying sequence of φ -modules is $(*)$. Remark that $p\mathfrak{N}' = 0$ and $p^{n-1}\mathfrak{N}'' = 0$. It is clear that $\hat{\mathfrak{N}}'' \in \text{Mod}_{\mathfrak{S}_\infty}^{r', \hat{G}, \text{cris}}$. Since

$$0 \rightarrow \widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}' \rightarrow \widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{N} \rightarrow \widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}'' \rightarrow 0$$

is exact and $p^{n-1}\mathfrak{N}'' = 0$, we obtain $\Delta(1 \otimes x) \in \widehat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}' \subset W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}'$ for any $x \in \mathfrak{M}$ by the induction hypothesis. Moreover, we have in fact $\Delta(1 \otimes x) \in u^p\varphi(\mathfrak{t})(W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}')$ since Lemma 7 implies

$$(W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}') \cap u^p\varphi(\mathfrak{t})(W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}) = u^p\varphi(\mathfrak{t})(W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}').$$

Identifying $W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}'$ with $R \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}'$, we obtain

$$\Delta(1 \otimes x) \in \mathfrak{m}_R^{\geq c(0)}(R \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}').$$

By an analogous argument of the case where $n = 1$, we obtain the implication

$$(s') \quad \text{For any } x \in \mathfrak{M}, \quad \Delta(1 \otimes x) \in \mathfrak{m}_R^{\geq c(s)}(R \otimes_{\varphi, \mathfrak{S}} \mathfrak{N}')$$

for any $s \geq 0$ and this implies $\Delta(1 \otimes x) = 0$. □

Before giving the proof of Theorem 6, we have to recall the theory of *maximal* Kisin modules. Now we give a very rough sketch of it (for more precise information, see [CL1]. Our sketch here is the case where “ $r = \infty$ ” in *loc. cit.*). For any $\mathfrak{M} \in \text{Mod}_{\mathfrak{S}_\infty}^r$, put $\mathfrak{M}[1/u] = \mathfrak{S}[1/u] \otimes_{\mathfrak{S}} \mathfrak{M}$ and denote by $F_{\mathfrak{S}}(\mathfrak{M}[1/u])$ the (partially) ordered set (by inclusion) of torsion Kisin modules \mathfrak{N} of finite height which is contained in $\mathfrak{M}[1/u]$ and $\mathfrak{N}[1/u] = \mathfrak{M}[1/u]$ as φ -modules. Here, a torsion Kisin module is called *of finite height* if it is of height $\leq s$ for some integer $s \geq 0$. The set $F_{\mathfrak{S}}(\mathfrak{M}[1/u])$ has a greatest element (cf. *loc. cit.*, Corollary 3.2.6), which is denoted by $\text{Max}(\mathfrak{M})$. We say that \mathfrak{M} is *maximal* if it is the greatest element of $F_{\mathfrak{S}}(\mathfrak{M}[1/u])$. The implication $\mathfrak{M} \mapsto \text{Max}(\mathfrak{M})$ defines a functor “Max” from the category of torsion Kisin modules of finite height into the category $\text{Max}_{/\mathfrak{S}_\infty}$ of maximal torsion Kisin modules. Furthermore, the functor $T_{\mathfrak{S}}: \text{Max}_{/\mathfrak{S}_\infty} \rightarrow \text{Rep}_{\text{tor}}(G_\infty)$, defined by $T_{\mathfrak{S}}(\mathfrak{M}) = \text{Hom}_{\mathfrak{S}, \varphi}(\mathfrak{M}, \mathbb{Q}_p/\mathbb{Z}_p \otimes_{\mathbb{Z}_p} W(R))$, is fully faithful (cf. *loc.*

cit., Corollary 3.3.10). It is not difficult to check that $T_{\mathfrak{S}}(\text{Max}(\mathfrak{M}))$ is canonically isomorphic to $T_{\mathfrak{S}}(\mathfrak{M})$ as representations of G_{∞} for any torsion Kisin module \mathfrak{M} .

Lemma 9. *Suppose $er < p - 1$. Then any $\mathfrak{M} \in \text{Mod}_{/\mathfrak{S}_{\infty}}^r$ is maximal.*

Proof. We prove by induction on n such that $p^n\mathfrak{M} = 0$. If $n = 1$, then the assertion follows by Lemma 3.3.4 of [CL1]. Suppose $n > 1$ and $p^n\mathfrak{M} = 0$. Take any $\mathfrak{N} \in F_{\mathfrak{S}}(\mathfrak{M}[1/u])$ such that $\mathfrak{M} \subset \mathfrak{N}$ and put $M = \mathfrak{M}[1/u] = \mathfrak{N}[1/u]$. Denote by pr the natural surjection $M \rightarrow M/pM$. Putting $\mathfrak{M}' = pM \cap \mathfrak{M}$, $\mathfrak{M}'' = \text{pr}(\mathfrak{M})$, $\mathfrak{N}' = pM \cap \mathfrak{N}$ and $\mathfrak{N}'' = \text{pr}(\mathfrak{N})$, we see that \mathfrak{M}' and \mathfrak{M}'' are objects of $\text{Mod}_{/\mathfrak{S}_{\infty}}^r$, and \mathfrak{N}' and \mathfrak{N}'' are torsion Kisin modules of finite height. Furthermore, we see that natural sequences

$$\begin{aligned} 0 \rightarrow \mathfrak{M}' \rightarrow \mathfrak{M} \xrightarrow{\text{pr}} \mathfrak{M}'' \rightarrow 0, \\ 0 \rightarrow \mathfrak{N}' \rightarrow \mathfrak{N} \xrightarrow{\text{pr}} \mathfrak{N}'' \rightarrow 0, \end{aligned}$$

of φ -modules are exact. By the induction hypothesis, we know that \mathfrak{M}' and \mathfrak{M}'' are maximal and thus $\mathfrak{N}' = \mathfrak{M}'$ and $\mathfrak{N}'' = \mathfrak{M}''$ (remark that $\mathfrak{M}'[1/u] = \mathfrak{N}'[1/u] = pM$ and $\mathfrak{M}''[1/u] = \mathfrak{N}''[1/u] = M/pM$). This implies $\mathfrak{N} = \mathfrak{M}$. \square

Proof of Theorem 6. Suppose that $er < p - 1$ and $e(r' - 1) < p - 1$. Let $T \in \text{Rep}_{\text{tor}}^{r, \hat{G}, \text{cris}}(G_K)$ (resp. $T' \in \text{Rep}_{\text{tor}}^{r', \hat{G}, \text{cris}}(G_K)$) and take $\hat{\mathfrak{M}} \in \text{Mod}_{/\mathfrak{S}_{\infty}}^{r, \hat{G}, \text{cris}}$ (resp. $\hat{\mathfrak{M}}' \in \text{Mod}_{/\mathfrak{S}_{\infty}}^{r', \hat{G}, \text{cris}}$) such that $T = \hat{T}(\hat{\mathfrak{M}})$ (resp. $T' = \hat{T}(\hat{\mathfrak{M}}')$). Note that $\mathfrak{M} = \text{Max}(\hat{\mathfrak{M}})$ by Lemma 9. By Theorem 3(1), we have the following commutative diagram:

$$\begin{array}{ccc} \text{Hom}_{G_K}(T, T') \hookrightarrow & & \text{Hom}_{G_{\infty}}(T, T') \\ \hat{T} \uparrow & & \uparrow T_{\mathfrak{S}} \\ \text{Hom}(\hat{\mathfrak{M}}', \hat{\mathfrak{M}}) \xrightarrow{\text{forgetful}} & \text{Hom}(\mathfrak{M}', \mathfrak{M}) \xrightarrow{\text{Max}} & \text{Hom}(\text{Max}(\mathfrak{M}'), \mathfrak{M}). \end{array}$$

The first bottom horizontal arrow is bijective by Lemma 8 and the second is also by an easy argument. Since the right vertical arrow is bijective, the top horizontal arrow must be bijective. \square

Remark 10. By Lemma 8, we can prove the latter part of Theorem 1 directly without using the former part of Theorem 1 as below: Suppose that $er < p - 1$. Let $T \in \text{Rep}_{\text{tor}}^r(G_K)$ (resp. $T' \in \text{Rep}_{\text{tor}}^{r'}(G_K)$) and take $\hat{\mathfrak{M}}$ (resp. $\hat{\mathfrak{M}}'$) be as in Theorem 4, which is an object of $\text{Mod}_{/\mathfrak{S}_{\infty}}^{r, \hat{G}, \text{cris}}$. By Theorem 3(1), we have the commutative diagram

$$\begin{array}{ccc} \text{Hom}_{G_K}(T, T') \hookrightarrow & & \text{Hom}_{G_{\infty}}(T, T') \\ \hat{T} \uparrow & & \uparrow T_{\mathfrak{S}} \\ \text{Hom}(\hat{\mathfrak{M}}', \hat{\mathfrak{M}}) \xrightarrow{\text{forgetful}} & & \text{Hom}(\mathfrak{M}', \mathfrak{M}) \end{array}$$

and then we obtain the desired result by an analogous argument before this remark.

Remark 11. The condition $e(r - 1) < p - 1$ in Lemma 8 is essential for the fullness of the forgetful functor $\text{Mod}_{/\mathfrak{S}_\infty}^{r, \hat{G}, \text{cris}} \rightarrow \text{Mod}_{/\mathfrak{S}_\infty}^r$ (note that this functor is always faithful). In fact, we have an example which implies that this forgetful functor is not full even if $e(r - 1) = p - 1$. We show below that the forgetful functor $\text{Mod}_{/\mathfrak{S}_\infty}^{r, \hat{G}, \text{cris}} \rightarrow \text{Mod}_{/\mathfrak{S}_\infty}^r$ is not full when $K = \mathbb{Q}_p$ and $r = p$.

Suppose $K = \mathbb{Q}_p$. Let E_π be the Tate curve over \mathbb{Q}_p associated with π . Lemma 18 in the next section says that the 2-dimensional \mathbb{F}_p -representation $E_\pi[p]$ of $G_{\mathbb{Q}_p}$ is torsion crystalline with Hodge–Tate weights in $[0, p]$. In particular, by Theorem 4, there exists a (φ, \hat{G}) -module $\hat{\mathfrak{M}} \in \text{Mod}_{/\mathfrak{S}_\infty}^{p, \hat{G}, \text{cris}}$ such that $\hat{T}(\hat{\mathfrak{M}}) \simeq E_\pi[p]$. On the other hand, for any nonnegative integer ℓ , define the (φ, \hat{G}) -module $\hat{\mathfrak{S}}_1(\ell) = (\mathfrak{S}_1(\ell), \varphi, \hat{G})$ as below: $\mathfrak{S}_1(\ell) = k[[u]] \cdot \mathfrak{f}^\ell$ is the rank-1 free $k[[u]]$ -module equipped with the Frobenius $\varphi(\mathfrak{f}^\ell) = c_0^{-\ell} u^{e\ell} \cdot \mathfrak{f}^\ell$, and define a \hat{G} -action on $\hat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{S}_1(\ell)$ by $\tau(\mathfrak{f}^\ell) = \hat{c}^\ell \cdot \mathfrak{f}^\ell$. Here, $\hat{c} = \prod_{n=1}^\infty \varphi^n(\frac{E(u)}{\tau(E(u))})$, which is contained in $\hat{\mathcal{R}}^\times$ (cf. Example 3.2.3 of [Li4]). Then Example 3.2.3 of *loc. cit.* says that $\hat{T}(\hat{\mathfrak{S}}_1(\ell)) \simeq \mathbb{F}_p(\ell)$. On the other hand, we define the (φ, \hat{G}) -module $\hat{\mathfrak{S}}_1(\ell)_0 = (\mathfrak{S}_1(\ell)_0, \varphi, \hat{G})$ as below: Put $\ell_0 = \max\{\ell' \in \mathbb{Z}_{\geq 0}; e\ell - (p-1)\ell' \geq 0\}$. We denote by $\mathfrak{S}_1(\ell)_0 = k[[u]] \cdot \mathfrak{g}^\ell$ the rank-1 free $k[[u]]$ -module equipped with the Frobenius $\varphi(\mathfrak{g}^\ell) = c_0^{-\ell} u^{e\ell - (p-1)\ell_0} \cdot \mathfrak{g}^\ell$, and define a \hat{G} -action on $\hat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{S}_1(\ell)$ by $\tau(\mathfrak{g}^\ell) = \underline{\varepsilon}^{-p\ell_0} \hat{c}^\ell \cdot \mathfrak{g}^\ell$. (The generator \mathfrak{g}^ℓ is taken to behave as $u^{-\ell_0} \mathfrak{f}^\ell$.) Then we see that $\text{Max}(\mathfrak{S}_1(\ell)) = \mathfrak{S}_1(\ell)_0$, and $\hat{\mathfrak{S}}_1(\ell)_0$ (and $\hat{\mathfrak{S}}_1(\ell)$) are objects of $\text{Mod}_{/\mathfrak{S}_\infty}^{\ell, \hat{G}, \text{cris}}$. We also see $\hat{T}(\hat{\mathfrak{S}}_1(\ell)_0) \simeq \hat{T}(\hat{\mathfrak{S}}_1(\ell)) \simeq \mathbb{F}_p(\ell)$. Now we consider the following commutative diagram (here we remark that $\mathfrak{S}_1(0) \oplus \mathfrak{S}_1(1)_0$ is maximal):

$$\begin{array}{ccc}
 \text{Hom}_{G_{\mathbb{Q}_p}}(\mathbb{F}_p \oplus \mathbb{F}_p(1), E_\pi[p]) & \hookrightarrow & \text{Hom}_{G_\infty}(\mathbb{F}_p \oplus \mathbb{F}_p(1), E_\pi[p]) \\
 \hat{T} \uparrow & & \uparrow T_{\mathfrak{S}} \\
 \text{Hom}(\hat{\mathfrak{M}}, \hat{\mathfrak{S}}_1(0) \oplus \hat{\mathfrak{S}}_1(1)_0) & & \\
 \downarrow \text{forgetful} & & \\
 \text{Hom}(\mathfrak{M}, \mathfrak{S}_1(0) \oplus \mathfrak{S}_1(1)_0) & \xrightarrow{\text{Max}} & \text{Hom}(\text{Max}(\mathfrak{M}), \mathfrak{S}_1(0) \oplus \mathfrak{S}_1(1)_0).
 \end{array}$$

The bottom horizontal arrow and the right vertical arrow are bijective since $\mathfrak{S}_1(0) \oplus \mathfrak{S}_1(1)_0$ is maximal. On the other hand, it is well-known that the inclusion $\text{Hom}_{G_{\mathbb{Q}_p}}(\mathbb{F}_p \oplus \mathbb{F}_p(1), E_\pi[p]) \subset \text{Hom}_{G_\infty}(\mathbb{F}_p \oplus \mathbb{F}_p(1), E_\pi[p])$ is not equal. Therefore,

$$\text{Hom}(\hat{\mathfrak{M}}, \hat{\mathfrak{S}}_1(0) \oplus \hat{\mathfrak{S}}_1(1)_0) \xrightarrow{\text{forgetful}} \text{Hom}(\mathfrak{M}, \mathfrak{S}_1(0) \oplus \mathfrak{S}_1(1)_0)$$

is not surjective. This implies that the forgetful functor

$$\text{Mod}_{\mathfrak{S}_\infty}^{p, \hat{G}, \text{cris}} \rightarrow \text{Mod}_{\mathfrak{S}_\infty}^p$$

is not full.

Remark 12. Combining Theorem (2.3.5) of [Kis], Theorem 4 and Lemma 8, we see that the forgetful functor $\text{Mod}_{\mathfrak{S}_\infty}^{1, \hat{G}, \text{cris}} \rightarrow \text{Mod}_{\mathfrak{S}_\infty}^1$ is an equivalence of categories.

4. Nonfullness: Examples

In the previous section, we showed that the restriction functor

$$\text{Rep}_{\text{tor}}^r(G_K) \xrightarrow{\text{res}} \text{Rep}_{\text{tor}}(G_\infty)$$

is fully faithful under the condition that $er < p - 1$. However, the full faithfulness may not hold if $er \geq p - 1$. In this section, we give some examples of this phenomenon. It should be noted that all our examples appearing in this section are given under the condition $e(r - 1) \geq p - 1$.

Let μ_{p^n} be the set of p^n -th roots of unity in \bar{K} , $\mu_{p^\infty} := \bigcup_{n \geq 0} \mu_{p^n}$ and denote by $G_1 \subset G_K$ the absolute Galois group of $K(\pi_1)$. Remark that, if the restriction functor $\mathcal{C} \rightarrow \text{Rep}_{\text{tor}}(G_1)$ is not fully faithful for a full subcategory \mathcal{C} of $\text{Rep}_{\text{tor}}(G_K)$, then the restriction functor $\mathcal{C} \rightarrow \text{Rep}_{\text{tor}}(G_\infty)$ is not fully faithful. Furthermore, we also remark that restriction functors $\mathcal{C} \rightarrow \text{Rep}_{\text{tor}}(G_\infty)$ and $\mathcal{C} \rightarrow \text{Rep}_{\text{tor}}(G_1)$ are always faithful.

Proposition 13. *Let K be a finite extension of \mathbb{Q}_p . Let s be the largest integer n such that $\mu_{p^n} \subset K$. Suppose that $s \geq 1$ and $K(\mu_{p^{s+1}})/K$ is ramified. Then the functor from torsion crystalline \mathbb{Z}_p -representations of G_K with Hodge–Tate weights in $[0, p + 1]$ to torsion \mathbb{Z}_p -representations of G_1 , obtained by restricting the action of G_K to G_1 , is not full.*

The lemma below follows from direct calculations.

Lemma 14. *Let $s \geq 1$ be an integer and $\psi: G_K \rightarrow \mathbb{Z}_p^\times$ an unramified character with the property that s is the largest integer n such that $\psi \bmod p^n$ is trivial. Define $\beta_\psi: G_K \rightarrow \mathbb{Z}_p$ by the relation $\psi = 1 + p^s \beta_\psi$ and put $\bar{\beta}_\psi = \beta \bmod p$. Denote by $\delta_\psi^0: H^0(G_K, \mathbb{Q}_p/\mathbb{Z}_p(\psi)) \rightarrow H^1(G_K, \mathbb{F}_p)$ the connection map coming from the exact sequence $0 \rightarrow \mathbb{F}_p \rightarrow \mathbb{Q}_p/\mathbb{Z}_p(\psi) \xrightarrow{p} \mathbb{Q}_p/\mathbb{Z}_p(\psi) \rightarrow 0$ of G_K -modules. Then $\bar{\beta}_\psi \in H^1(G_K, \mathbb{F}_p)$ and $\text{Im}(\delta_\psi^0) = \mathbb{F}_p \cdot \bar{\beta}_\psi$.*

Proof of Proposition 13. Let $\varepsilon: G_K \rightarrow \mathbb{Z}_p^\times$ be the p -adic cyclotomic character and $\bar{\varepsilon} := \varepsilon \bmod p$ the mod p cyclotomic character. Let K and $s \geq 1$ be as in Proposition 13. Let $\chi: G_K \rightarrow \mathbb{Z}_p^\times$ be an unramified character such that $\chi \bmod p^s$ is trivial. It suffices to show that, for some choice of χ , there exist $\rho: G_K \rightarrow GL_2(\mathbb{Z}_p)$ and $2 \leq r \leq p + 1$ with an exact sequence $0 \rightarrow \chi \varepsilon^r \rightarrow \rho \rightarrow 1 \rightarrow 0$ of representations of G_K such that $\rho \bmod p$ is not trivial on G_K but is trivial on G_1 . Here, 1 in the

above exact sequence means the trivial character. Note that such ρ is always crystalline (cf. [BK, Example 3.9]). Since $\mu_p \subset K$, we can define $f_0 \in H^1(G_K, \mathbb{F}_p)$ such that f_0 factors through \hat{G} , $f_0(\tau) = 1$ and $f_0|_{H_K} = 0$, where H_K is defined in Definition 2. The kernel of the restriction map $H^1(G_K, \mathbb{F}_p) \rightarrow H^1(G_1, \mathbb{F}_p)$ is a one dimensional \mathbb{F}_p -vector space which is generated by f_0 . Let $H \subset H^1(G_K, \mathbb{F}_p)$ be an annihilator of f_0 under the Tate paring. For any integer ℓ , denote by $\delta_{\chi, \ell}^1: H^1(G_K, \mathbb{F}_p) \rightarrow H^2(G_K, \mathbb{Z}_p(\chi\varepsilon^\ell))$ (resp. $\delta_{\chi, \ell}^0: H^0(G_K, \mathbb{Q}_p/\mathbb{Z}_p(\chi^{-1}\varepsilon^{1-\ell})) \rightarrow H^1(G_K, \mathbb{F}_p)$) the connection map coming from the exact sequence $0 \rightarrow \mathbb{Z}_p(\chi\varepsilon^\ell) \xrightarrow{p} \mathbb{Z}_p(\chi\varepsilon^\ell) \rightarrow \mathbb{F}_p \rightarrow 0$ (resp. $0 \rightarrow \mathbb{F}_p \rightarrow \mathbb{Q}_p/\mathbb{Z}_p(\chi^{-1}\varepsilon^{1-\ell}) \xrightarrow{p} \mathbb{Q}_p/\mathbb{Z}_p(\chi^{-1}\varepsilon^{1-\ell}) \rightarrow 0$) of G_K -modules. By Tate local duality, the condition that f_0 lifts to $H^1(G_K, \mathbb{Z}_p(\chi\varepsilon^\ell))$ is equivalent to the condition that $\text{Im}(\delta_{\chi, \ell}^0) \subset H$. Hence it is enough to choose χ which satisfies the latter condition for some $2 \leq \ell \leq p + 1$.

Since $K(\mu_{p^{s+1}})/K$ is ramified, we know that s is the largest integer n such that $\chi^{-1}\varepsilon^{-1} \bmod p^n$ is trivial. Take $\beta_{\chi^{-1}\varepsilon^{-1}}$ and $\bar{\beta}_{\chi^{-1}\varepsilon^{-1}}$ as in Lemma 14. For simplicity, we write $\alpha_\chi := \beta_{\chi^{-1}\varepsilon^{-1}}$ and $\bar{\alpha}_\chi := \bar{\beta}_{\chi^{-1}\varepsilon^{-1}}$. By Lemma 14, $\text{Im}(\delta_{\chi, 2}^0)$ is generated by $\bar{\alpha}_\chi$. If $\bar{\alpha}_1$ is contained in H , then we finish the proof (choose χ as the trivial character 1). Suppose $\bar{\alpha}_1$ is not contained in H . From now on, we fix χ as follows; χ is the unramified character $G_K \rightarrow \mathbb{Z}_p^\times$ with $\chi(\text{Frob}_K) = (1 + p^s)^{-1}$, where Frob_K is the arithmetic Frobenius of K . Let $u_1: G_K \rightarrow \mathbb{F}_p$ be the unramified homomorphism with $u_1(\text{Frob}_K) = 1$. Then we obtain $\bar{\alpha}_\chi = u_1 + \bar{\alpha}_1$. Since $K(\mu_{p^{s+1}})/K$ is ramified, we see that $\bar{\alpha}_1|_{I_K}$ is not zero where I_K is the inertia subgroup of G_K . This implies $u_1 \notin \mathbb{F}_p \cdot \bar{\alpha}_1$. Noting that $H^1(G_K, \mathbb{F}_p) = H \oplus \mathbb{F}_p \cdot \bar{\alpha}_1$, we have $\bar{\alpha}_\chi + \bar{a}\bar{\alpha}_1 \in H$ for some $\bar{a} \in \mathbb{F}_p$. Let $0 \leq a \leq p - 1$ be the integer such that $a \bmod p$ is \bar{a} . Under the modulo p^{2s} , we have $\chi^{-1}\varepsilon^{-(1+a)} = \chi^{-1}\varepsilon^{-1} \cdot \varepsilon^{-a} = (1 + p^s\alpha_\chi)(1 + p^sa\alpha_1) = 1 + p^s(\alpha_\chi + a\alpha_1)$. Since $\bar{\alpha}_\chi + \bar{a}\bar{\alpha}_1 = u_1 + (\bar{a} + 1)\bar{\alpha}_1 \neq 0$, we see that s is the largest integer n such that $\chi^{-1}\varepsilon^{-(1+a)} \bmod p^n$ is trivial. Hence, defining $\bar{\beta}_{\chi^{-1}\varepsilon^{-(1+a)}}$ as in Lemma 14, we obtain $\bar{\beta}_{\chi^{-1}\varepsilon^{-(1+a)}} = \bar{\alpha}_\chi + \bar{a}\bar{\alpha}_1$. Therefore, we obtain that $\text{Im}(\delta_{\chi, 2+a}^0) = \mathbb{F}_p \cdot \bar{\beta}_{\chi^{-1}\varepsilon^{-(1+a)}} \subset H$ and we are done. \square

Unfortunately, Proposition 13 can not be applied even when $K = \mathbb{Q}_p$. On the other hand, the following proposition is effective for $K = \mathbb{Q}_p$, but we need a certain restriction on the choice of the uniformizer π . Let L be the unique degree p extension of K which is contained in $K(\mu_{p^\infty})$.

Proposition 15. *Let K be a finite extension of \mathbb{Q}_p . Suppose that π is contained in $\text{Norm}_{L/K}(L^\times)$. (Thus the extension L/K must be totally ramified in this case.) Then the functor from torsion crystalline \mathbb{Z}_p -representations of G_K with Hodge–Tate weights in $[0, p]$ to torsion \mathbb{Z}_p -representations of G_1 , obtained by restricting the action of G_K to G_1 , is not full.*

Proof. Let s be the largest integer n such that $\mu_{p^n} \subset K$. Then we can write $\varepsilon^{1-p} = 1 + p\psi$ with some map $\psi: G_K \rightarrow \mathbb{Z}_p$. Putting $\bar{\psi} = \psi \bmod p: G_K \rightarrow$

\mathbb{F}_p , we see that $\bar{\psi}$ is nontrivial homomorphism with kernel $\text{Gal}(\bar{K}/L)$. Let $\delta^0: H^0(G_K, \mathbb{Q}_p/\mathbb{Z}_p(1-p)) \rightarrow H^1(G_K, \mathbb{F}_p)$ be the connection map arising from the exact sequence $0 \rightarrow \mathbb{F}_p \rightarrow \mathbb{Q}_p/\mathbb{Z}_p(1-p) \xrightarrow{p} \mathbb{Q}_p/\mathbb{Z}_p(1-p) \rightarrow 0$. Under the isomorphism $K^\times/(K^\times)^p \simeq H^1(G_K, \mathbb{F}_p(1))$ via Kummer theory, $\pi \bmod (K^\times)^p$ corresponds to the 1-cocycle $[\pi]$ defined by $\sigma \mapsto \frac{\sigma(\pi_1)}{\pi_1}$ for $\sigma \in G_K$, which is clearly trivial on G_1 . By Tate local duality and the fact that the image of δ^0 is generated by $\bar{\psi}$ (cf. Lemma 14), it suffices to show that $([\pi], \bar{\psi})$ maps to zero under the Tate pairing $H^1(G_K, \mathbb{F}_p(1)) \times H^1(G_K, \mathbb{F}_p) \rightarrow \mathbb{Q}/\mathbb{Z}$ (in fact, this implies that $[\pi]$ lifts to $H^1(G_K, \mathbb{Z}_p(p))$ and we obtain the desired result). Let $\phi_{L/K}: K^\times/\text{Norm}_{L/K}(L^\times) \xrightarrow{\sim} \text{Gal}(L/K)$ be the isomorphism of local class field theory. It is enough to show that $\bar{\psi}(\phi_{L/K}(\pi)) = 0$. Our assumption of π implies that this equality certainly holds. \square

Now we give an example for the nonfullness of our restriction functor without any assumption on the choice of the uniformizer π .

Proposition 16. *The functor from torsion crystalline \mathbb{Z}_p -representations of $G_{\mathbb{Q}_p}$ with Hodge–Tate weights in $[0, p]$ to torsion \mathbb{Z}_p -representations of G_1 , obtained by restricting the action of $G_{\mathbb{Q}_p}$ to G_1 , is not full.*

Lemma 17. *Let F be a finite extension of \mathbb{Q}_p or \mathbb{F}_p . Then any 2-dimensional irreducible F -representation of $G_{\mathbb{Q}}$ whose determinant is the p -adic cyclotomic character is absolutely irreducible.*

Proof. Let $\rho: G_{\mathbb{Q}} \rightarrow GL_2(F)$ be as in the statement and denote by V the underlying F -vector space. Suppose that, for some finite extension F' over F , there exists a $G_{\mathbb{Q}}$ -stable F' -subvector space W of $F' \otimes_F V$ of dimension 1. If we denote by $c \in G_{\mathbb{Q}}$ the complex conjugation, then $\rho(c)^2$ is the identity matrix and $\det(\rho)(c) = -1$. Hence it follows that $\rho(c)$ is conjugate (over F) with $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ (note that p is odd). By this fact and the fact that $\rho(c)$ preserves W , we see that W is defined over F . This is a contradiction. \square

Lemma 18. *Let K be a finite extension of \mathbb{Q}_p and $q \in \mathbb{Q}_p^\times (K^\times)^p$. Let $E_q[p]$ be the Tate curve over K associated with q . If $p \nmid v_K(q)$, then $E_q[p]$ is torsion crystalline with Hodge–Tate weights in $[0, p]$.*

Proof. We have a decomposition $q = q'q''$, where $q' \in \mathbb{Q}_p^\times, v_K(q') > 0$ and $q'' \in (K^\times)^p$. Let $E_{q'}$ be the Tate curve over \mathbb{Q}_p associated with q' . Then $E_{q'}[p]$ is a representation of $G_{\mathbb{Q}_p}$ and we have an isomorphism $E_q[p] \simeq (E_{q'}[p])|_{G_K}$. Hence we can reduce the case where $K = \mathbb{Q}_p$. Let $\ell > 3$ be a prime number different from p such that $-\ell$ is not a square in \mathbb{F}_p^\times (recall that p is odd). Choose an elliptic curve $E_{(\ell)}$ over \mathbb{Q}_ℓ which has good supersingular reduction. Since $\ell > 3$, we have $\#E_{(\ell)}(\mathbb{F}_\ell) = 1 + \ell$. Thus the characteristic polynomial of $E_{(\ell)}[p]$ for the arithmetic Frobenius of ℓ is $X^2 + \ell \in \mathbb{F}_p[X]$, which does not have a root in \mathbb{F}_p . Hence $E_{(\ell)}[p]$ is an irreducible representation of $G_{\mathbb{Q}_\ell}$ where $G_{\mathbb{Q}_\ell}$ is the absolute Galois group of

\mathbb{Q}_ℓ . We define \mathcal{S} to be the set of \mathbb{Q} -isomorphism classes of elliptic curves E defined over \mathbb{Q} which satisfy the following conditions:

- (a) E has multiplicative reduction at p and

$$v_p(j(E)) = v_p(j(E_q)) (= -v_p(q))$$

where $j(E)$ is the j -invariant of E .

- (b) $E[p] \simeq E_q[p]$ as \mathbb{F}_p -representations of $G_{\mathbb{Q}_p}$.
- (c) $E[p] \simeq E_{(\ell)}[p]$ as \mathbb{F}_p -representations of $G_{\mathbb{Q}_\ell}$.

The set \mathcal{S} is infinite since elliptic curves over \mathbb{Q} , whose coefficients of their defining equations are p -adically close enough to that of E_π and also ℓ -adically close enough to that of $E_{(\ell)}$, are contained in \mathcal{S} . Now we take any elliptic curve E over \mathbb{Q} whose \mathbb{Q} -isomorphism class is in the set \mathcal{S} . By the condition (c), $E[p]$ is irreducible as a representation of $G_{\mathbb{Q}}$. It is moreover absolutely irreducible by Lemma 17. By the classical Serre’s modularity conjecture (proved by Khare and Wintenberger) and the well-known fact that p -adic representations arising from Hecke eigencusp forms of level prime to p are crystalline, we know that $(E[p] \otimes_{\mathbb{F}_p} \overline{\mathbb{F}_p})|_{G_{\mathbb{Q}_p}}$ is the reduction of a lattice in some crystalline $\overline{\mathbb{Q}_p}$ -representation. Furthermore, by the condition (a) and Proposition 5(2) of [Se], we know that $(E[p])|_{G_{\mathbb{Q}_p}}$ is torsion crystalline with Hodge–Tate weights in $[0, p]$. Therefore, so is $E_q[p]$ by (b). \square

Proof of Proposition 16. Put $T = E_\pi[p]$ and $T' = \mathbb{F}_p \oplus \mathbb{F}_p(1)$. We know that T and T' are in $\text{Rep}_{\text{tor}}^p(G_{\mathbb{Q}_p})$ by Lemma 18. They are not isomorphic as representations of $G_{\mathbb{Q}_p}$ but isomorphic as representations of G_1 . This gives the desired result. \square

Here we suggest the following question.

Question 19. *What is the necessary and sufficient condition for that the functor*

$$\text{Rep}_{\text{tor}}^r(G_K) \xrightarrow{\text{res}} \text{Rep}_{\text{tor}}(G_\infty), \quad T \mapsto T|_{G_\infty}$$

is fully faithful? Furthermore, does this condition depend only on e and r ?

Remark 20.

- (1) We do not know whether the full faithfulness of the functor in Question 19 depends on the choice of the system $(\pi_n)_{n \geq 0}$ or not (see Proposition 15). However, it is not difficult to see the following: Take two systems $(\pi_n)_{n \geq 0}$ and $(\pi'_n)_{n \geq 0}$ of p^n -th roots of a fixed uniformizer π of K (thus we have $\pi_0 = \pi'_0 = \pi$). Put $K_\infty = \bigcup_{n \geq 0} K(\pi_n)$ (resp. $K'_\infty = \bigcup_{n \geq 0} K(\pi'_n)$) and $G_\infty = \text{Gal}(\overline{K}/K_\infty)$ (resp. $G'_\infty = \text{Gal}(\overline{K}/K'_\infty)$). Then, the restriction functor

$$\text{Rep}_{\text{tor}}^r(G_K) \xrightarrow{\text{res}} \text{Rep}_{\text{tor}}(G_\infty)$$

is fully faithful if and only if the restriction functor

$$\text{Rep}_{\text{tor}}^r(G_K) \xrightarrow{\text{res}} \text{Rep}_{\text{tor}}(G'_\infty)$$

is. In fact, we can check this from the fact that G_∞ and G'_∞ are conjugate with each other by some element of G_K .

- (2) A torsion \mathbb{Z}_p -representation of G_K is called *finite flat* if it is isomorphic to $G(\overline{K})$ as \mathbb{Z}_p -representations of G_K for some p -power order finite flat commutative group scheme G over the integer ring of K . If $r = 1$, then the category $\text{Rep}_{\text{tor}}^r(G_K) = \text{Rep}_{\text{tor}}^1(G_K)$ coincides with the category of finite flat representations of G_K (this can be checked by, for example, Theorem 3.1.1 of [BBM]). Breuil proved in Theorem 3.4.3 of [Br3] that the restriction functor

$$\text{Rep}_{\text{tor}}^1(G_K) \xrightarrow{\text{res}} \text{Rep}_{\text{tor}}(G_\infty)$$

is fully faithful for any K without any restriction on e . In fact, this assertion is true even if $p = 2$ (cf. [Kim], [La], [Li4], proved independently. Explicitly, see Corollary 4.4 of [Kim]).

- (3) If $e = 1$ and $r < p - 1$, then the fact that the restriction functor $\text{Rep}_{\text{tor}}^r(G_K) \xrightarrow{\text{res}} \text{Rep}_{\text{tor}}(G_\infty)$ is fully faithful has been already known ([Br2], the proof of Théorème 5.2).
- (4) Observing known results as above and results shown in this paper, it seems that the answer of Question 19 should be “ $e(r - 1) < p - 1$ ”.

Appendix A. (φ, \hat{G}) -modules associated with crystalline representations

In Proposition 5.9 of [GLS], a necessary condition for representations arising from free (φ, \hat{G}) -modules to be crystalline is given. In this appendix, we show that the converse holds. The result here justifies the subscript “cris” of the category $\text{Mod}_{\mathfrak{S}_\infty}^{r, \hat{G}, \text{cris}}$ defined in Section 3.

We continue to use the same notation as in Section 2. For any integer $n \geq 0$, we define ideals of $W(R)$ as below:

$$I^{[n]}W(R) := \{a \in W(R); \varphi^m(a) \in \text{Fil}^n A_{\text{cris}} \text{ for every } m \geq 0\},$$

$$I^{[n^+]}W(R) := I^{[n]}W(R)I_+W(R)$$

(see Section 5 of [Fo2] for more precise information). The proof of Lemma 3.2.2 of [Li2] shows that $I^{[n]}W(R)$ is a principal ideal of $W(R)$ generated by $\varphi(\mathfrak{t})^n$. In particular we see that $u^p \varphi(\mathfrak{t})$ is contained in

$$I^{[1^+]}W(R) = \varphi(\mathfrak{t})I_+W(R).$$

Recall that $\hat{T}(\hat{\mathfrak{M}}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ is a semi-stable \mathbb{Q}_p -representation of G_K (Theorem 3(2)) and $\tau(x) - x \in I_+W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$ for any $x \in \mathfrak{M}$. The main purpose of this appendix is to prove the following:

Theorem 21. *Let $\hat{\mathfrak{M}} \in \text{Mod}_{\mathfrak{S}}^{r, \hat{G}}$ be a (φ, \hat{G}) -module. The following are equivalent:*

- (1) $\hat{T}(\hat{\mathfrak{M}}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ is crystalline.

- (2) For any $x \in \mathfrak{M}$, we have $\tau(x) - x \in I^{[1^+]}W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$.
- (3) For any $x \in \mathfrak{M}$, we have $\tau(x) - x \in u^p \varphi(\mathfrak{t})(W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M})$.

Before giving a proof of this theorem, we shall recall some known facts about (φ, \hat{G}) -modules. Let $\hat{\mathfrak{M}} \in \text{Mod}_{\mathfrak{S}}^{r, \hat{G}}$ be a (φ, \hat{G}) -module, and put $\mathcal{D} = S_{K_0} \otimes_{\varphi, \mathfrak{S}} \hat{\mathfrak{M}}$. Then \mathcal{D} has a structure as a Breuil module which corresponds to the semi-stable representation $\hat{T}(\hat{\mathfrak{M}}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ of G_K . (Breuil modules here are objects of “ $\mathcal{MF}_S(\varphi, N)$ ” defined in Section 6.1 of [Br1]. It is useful for the reader to refer also Section 5 of [Li1].) Denote by $N_{\mathcal{D}}$ the monodromy operator of \mathcal{D} and define a G_K -action on $B_{\text{cris}}^+ \otimes_S \mathcal{D} = B_{\text{cris}}^+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$ by

$$g(a \otimes x) = \sum_{i=0}^{\infty} g(a) \gamma_i(-\log([\underline{\varepsilon}])) \otimes N_{\mathcal{D}}(x)$$

for $g \in G_K, a \in B_{\text{cris}}^+, x \in \mathcal{D}$. By the construction of the quasi-inverse of the functor \hat{T} of Theorem 3(2) ([Li2], Section 3.2), this G_K -action is stable on $\hat{\mathcal{R}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M} \subset B_{\text{cris}}^+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$ and it factors through \hat{G} , which gives the original \hat{G} -action of the (φ, \hat{G}) -module $\hat{\mathfrak{M}}$. For any $n \geq 0$ and any $x \in \mathcal{D}$, an induction on n shows that

$$(\tau - 1)^n(x) = \sum_{m=n}^{\infty} \left\{ \sum_{i_1 + \dots + i_n = m, i_j \geq 0} \frac{m!}{i_1! \dots i_n!} \right\} \gamma_m(t) \otimes N_{\mathcal{D}}^m(x) \in B_{\text{cris}}^+ \otimes_S \mathcal{D}$$

and in particular $\frac{(\tau - 1)^n}{n}(x) \rightarrow 0$ p -adically as $n \rightarrow \infty$. Hence we can define

$$\log(\tau)(x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{(\tau - 1)^n}{n}(x) \in B_{\text{cris}}^+ \otimes_S \mathcal{D}.$$

It is not difficult to check the equation $\log(\tau)(x) = t \otimes N_{\mathcal{D}}(x)$. Consequently the monodromy operator $N_{\mathcal{D}}$ can be reconstructed from the τ -action of $\hat{\mathfrak{M}}$ by the relation $\frac{1}{t} \log(\tau)(x) = N_{\mathcal{D}}(x)$. Put $D = \mathcal{D}/I_+ S_{K_0} \mathcal{D}$. Then D has a structure as a filtered (φ, N) -module over K_0 which corresponds to $\hat{T}(\hat{\mathfrak{M}}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ and the monodromy operator N_D of D is given by $N_{\mathcal{D}}$ mod $I_+ S_{K_0} \mathcal{D}$ ([Br1], Section 6). Hence $\hat{T}(\hat{\mathfrak{M}}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ is crystalline if and only if $N_{\mathcal{D}}$ mod $I_+ S_{K_0} \mathcal{D}$ is zero.

Proof of Theorem 21. The implication (1) \Rightarrow (3) follows from Proposition 5.9 of [GLS]. It is clear that (3) implies (2). Thus it suffices to show the implication (2) \Rightarrow (1). Assume the condition (2). We use the same notation $\mathcal{D}, N_{\mathcal{D}}, D, N_D$ as the above. We often regard \mathfrak{M} as a $\varphi(\mathfrak{S})$ -submodule of \mathcal{D} . Let $x \in \mathfrak{M}$. For any integer $n > 0$, it is shown in the proof of Proposition 2.4.1 of [Li3] that:

(A) $(\tau - 1)^n(x) \in I^{[n]}W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$.

(B) $\frac{(\tau-1)^n}{nt}(x)$ is well-defined in $A_{\text{cris}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$ and $\frac{(\tau-1)^n}{nt}(x) \rightarrow 0$ p -adically as $n \rightarrow \infty$. Therefore, we have

$$\frac{1}{t} \log(\tau)(x) \in A_{\text{cris}} \otimes_{\varphi, \mathfrak{S}} \mathfrak{M} \subset B_{\text{cris}}^+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}.$$

By (A), we can take $y_n \in W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$ such that $(\tau - 1)^n(x) = \varphi(\mathfrak{t})^n y_n$. Then we have the equation

$$(*) \quad cN_{\mathcal{D}}(x) = c \cdot \frac{1}{t} \log(\tau)(x) = \frac{\tau - 1}{\varphi(\mathfrak{t})}(x) + \sum_{n=2}^{\infty} (-1)^{n-1} \frac{\varphi(\mathfrak{t})^{n-1}}{n} y_n.$$

Here $c = \frac{t}{\varphi(\mathfrak{t})}$, which is a unit of A_{cris} ([Li2], Example 3.2.3). Note that $\frac{\tau-1}{\varphi(\mathfrak{t})}(x)$ is contained in $I_+W(R) \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$ by the assumption (2).

Now we claim that there exists an integer $n_0 > 1$ such that $\frac{(n-2)!}{n}$ is in \mathbb{Z}_p for any $n > n_0$. Admitting this claim, we proceed a proof of Theorem 21. Consider the decomposition

$$\begin{aligned} \sum_{n=2}^{\infty} (-1)^{n-1} \frac{\varphi(\mathfrak{t})^{n-1}}{n} y_n &= \\ \varphi(\mathfrak{t}) \sum_{n=2}^{n_0} (-1)^{n-1} \frac{\varphi(\mathfrak{t})^{n-2}}{n} y_n &+ \varphi(\mathfrak{t}) \sum_{n=n_0+1}^{\infty} (-1)^{n-1} \frac{\varphi(\mathfrak{t})^{n-2}}{n} y_n. \end{aligned}$$

By the claim, we see that $\frac{\varphi(\mathfrak{t})^{n-2}}{n} = \frac{\varphi(\mathfrak{t})^{n-2}}{(n-2)!} \cdot \frac{(n-2)!}{n} = c^{-(n-2)} \gamma_{n-2}(t) \frac{(n-2)!}{n}$ is contained in A_{cris} for any $n > n_0$ and it goes to zero p -adically as $n \rightarrow \infty$. In particular, (the first term and) the second term of the above decomposition are contained in $\varphi(\mathfrak{t})(B_{\text{cris}}^+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{M})$, which is contained in $I_+B_{\text{cris}}^+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$. Hence $\sum_{n=2}^{\infty} (-1)^{n-1} \frac{\varphi(\mathfrak{t})^{n-1}}{n} y_n$ is also contained in $I_+B_{\text{cris}}^+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$. Note that $\nu(c) = 1$ since $c = \frac{t}{\varphi(\mathfrak{t})} = \prod_{n=0}^{\infty} \varphi^n(\frac{c_0^{-1}E(u)}{p})$ and $\nu(u) = 0$, and furthermore $\nu(\mathfrak{t}) = 0$. Therefore, by (*) modulo $I_+B_{\text{cris}}^+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{M}$, we obtain the relation $N_D(\bar{x}) = 0$ in $D = \mathcal{D}/I_+S_{K_0}\mathcal{D} \subset (B_{\text{cris}}^+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{M})/(I_+B_{\text{cris}}^+ \otimes_{\varphi, \mathfrak{S}} \mathfrak{M})$ where \bar{x} is the residue class of x . Since the image of \mathfrak{M} in $D = \mathcal{D}/I_+S_{K_0}\mathcal{D}$ generates D as a K_0 -vector space, we obtain that $N_D = 0$. This implies (1). Hence it suffices to show the claim. Let v_p be the p -adic valuation with $v_p(p) = 1$. For any positive integer n , write $n = p^s m$ with $p \nmid m$. If $s = 0$, it is clear that $\frac{(n-2)!}{n} \in \mathbb{Z}_p$. Suppose $s \geq 1$. If $m \geq 2$, we have $v_p((n-2)!) \geq v_p((2p^s - 2)!) \geq v_p(p^s!) \geq s = v_p(n)$. If $m = 1$ and $s \geq 3$, we have

$$v_p((n - 2)!) \geq v_p(p^{s-1}!) = \frac{1}{2}s(s - 1) \geq s = v_p(n).$$

This finishes the proof. □

References

- [BBM] BERTHELOT, PIERRE; BREEN, LAWRENCE; MESSING, WILLIAM. Théorie de Dieudonné cristalline. II. Lecture Notes in Mathematics, 930. *Springer-Verlag, Berlin*, 1982. x+261 pp. ISBN: 3-540-11556-0. [MR0667344](#) (85k:14023), [Zbl 0516.14015](#), doi: [10.1007/BFb0093025](#).
- [BK] BLOCH, SPENCER; KATO, KAZUYA. L -functions and Tamagawa numbers of motives. *The Grothendieck Festschrift, I* 333–400, Progr. Math., 86. *Birkhäuser Boston, Boston, MA*, 1990. [MR1086888](#) (92g:11063), [Zbl 0768.14001](#), doi: [10.1007/978-0-8176-4574-8_9](#).
- [Br1] BREUIL, CHRISTOPHE. Représentations p -adiques semi-stables et transversalité de Griffiths. *Math. Ann.* **307** (1997), no. 2, 191–224. [MR1428871](#) (98b:14016), [Zbl 0883.11049](#), doi: [10.1007/s002080050031](#).
- [Br2] BREUIL, CHRISTOPHE. Une application de corps des normes. *Compositio Math.* **117** (1999) no. 2, 189–203. [MR1695849](#) (2000f:11157), [Zbl 0933.11055](#), doi: [10.1023/A:1000923331053](#).
- [Br3] BREUIL, CHRISTOPHE. Integral p -adic Hodge theory. *Algebraic geometry 2000, Azumino (Hotaka)*, 51–80, Adv. Stud. Pure Math., 36, *Math. Soc. Japan, Tokyo*, 2002. [MR1971512](#) (2004e:11135), [Zbl 1046.11085](#).
- [CL1] CARUSO, XAVIER; LIU, TONG. Quasi-semi-stable representations. *Bull. Soc. Math. France* **137** (2009), no. 2, 185–223. [MR2543474](#) (2011c:11086), [Zbl 1266.11073](#), [arXiv:0709.2118](#).
- [CL2] CARUSO, XAVIER; LIU, TONG. Some bounds for ramification of p^n -torsion semi-stable representations. *J. Algebra* **325** (2011), 70–96. [MR2745530](#) (2012b:11090), [Zbl 1269.14001](#), [arXiv:0805.4227](#), doi: [10.1016/j.jalgebra.2010.10.005](#).
- [Fo1] FONTAINE, JEAN-MARC. Représentations p -adiques des corps locaux. I. *The Grothendieck Festschrift, Vol. II*, 249–309, Progr. Math., 87. *Birkhäuser Boston, Boston, MA*, 1990. [MR1106901](#) (92i:11125), [Zbl 0743.11066](#).
- [Fo2] FONTAINE, JEAN-MARC. Le corps des périodes p -adiques. *Astérisque* **223** (1994), 59–111. [MR1293971](#) (95k:11086), [Zbl 0940.14012](#).
- [GLS] GEE, TOBY; LIU, TONG; SAVITT, DAVID. The Buzzard–Diamond–Jarvis conjecture for unitary groups. *J. Amer. Math. Soc.* **27** (2014), no. 2, 389–435. [MR3164985](#), [Zbl 1288.11045](#), [arXiv:1203.2552](#), doi: [10.1090/S0894-0347-2013-00775-4](#).
- [Kim] KIM, WANSU. The classification of p -divisible groups over 2-adic discrete valuation rings. *Math. Res. Lett.* **19** (2012), no. 1, 121–141. [MR2923180](#), [Zbl 1284.14056](#), [arXiv:1007.1904](#), doi: [10.4310/MRL.2012.v19.n1.a10](#).
- [Kis] KISIN, MARK. Crystalline representations and F -crystals. *Algebraic geometry and number theory*, 459–496, Progr. Math., 253. *Birkhäuser Boston, Boston, MA*, 2006. [MR2263197](#) (2007j:11163), [Zbl 1184.11052](#), doi: [10.1007/978-0-8176-4532-8_7](#).
- [La] LAU, EIKE. A relation between Dieudonné displays and crystalline Dieudonné theory. 2010. [arXiv:1006.2720](#).
- [Li1] LIU, TONG. Torsion p -adic Galois representations and a conjecture of Fontaine. *Ann. Sci. École Norm. Sup. (4)* **40** (2007), no. 4, 633–674. [MR2191528](#) (2010h:11191), [Zbl 1163.11043](#), doi: [10.1016/j.ansens.2007.05.002](#).
- [Li2] LIU, TONG. A note on lattices in semi-stable representations. *Math. Ann.* **346** (2010), no. 1, 117–138. [MR2558890](#) (2011d:11272), [Zbl 1208.14017](#), doi: [10.1007/s00208-009-0392-y](#).
- [Li3] LIU, TONG. Lattices in filtered (φ, N) -modules. *J. Inst. Math. Jussieu* **11**, (2012), no. 3, 659–693. [MR2931320](#), [Zbl 1249.14006](#), doi: [10.1017/S1474748011000235](#).

- [Li4] LIU, TONG. The correspondence between Barsotti–Tate groups and Kisin modules when $p = 2$. *J. Théor. Nombres Bordeaux* **25** (2013), no. 3, 661–676. [MR3179680](#), [Zbl 06291371](#), doi: [10.5802/jtnb.852](#).
- [Oz] OZEKI, YOSHIYASU. Torsion representations arising from (φ, \hat{G}) -modules. *J. Number Theory* **133** (2013), no. 11, 3810–3861. [MR3084302](#), [Zbl 1297.11146](#), [arXiv:1202.1858](#), doi: [10.1016/j.jnt.2013.05.010](#).
- [Se] SERRE, JEAN–PIERRE. Sur les représentations modulaires de degré 2 de $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$. *Duke Math. J* **54** (1987), no. 1, 179–230. [MR0885783](#) (88g:11022), [Zbl 0641.10026](#), doi: [10.1215/S0012-7094-87-05413-5](#).

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN.

yozeaki@kurims.kyoto-u.ac.jp

This paper is available via <http://nyjm.albany.edu/j/2014/20-51.html>.