

ON CLOSE-TO-CONVEX FUNCTIONS

D. K. Thomas

Abstract. Well-known coefficient and length results for the class of univalent close-to-convex functions are extended to a subclass of close-to-convex functions of high order.

1. Introduction. In [3] Goodman introduced the class $K(\beta)$ of normalised analytic functions which are close-to-convex of order $\beta \geq 0$, i.e. $f \in K(\beta)$ if f is analytic in $D = \{z : |z| < 1\}$ and if there exists $\varphi \in K(0) = C$ the class of normalised convex functions, such that for $z \in D$,

$$\left| \arg \frac{f'(z)}{\varphi'(z)} \right| \leq \frac{\beta\pi}{2}.$$

When $0 \leq \beta \leq 1$, $K(\beta)$ consists of univalent functions, whilst if $\beta > 1$ f need not even be finitely valent.

Denote by V_k , ($k \geq 2$) the class of locally univalent functions with bounded boundary rotation and by R_k the class of functions with bounded radial rotation. Then $\varphi \in V_k$ if, and only if, $z\varphi' \in R_k$ (see e.g. [2]). In [5] Noor considered the class T_k defined as follows:

Definition. Let $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ be analytic and locally univalent in D . Then for $k \geq 2$, $f \in T_k$ if there is a function $\varphi \in V_k$ such that for $z \in D$,

$$\operatorname{Re} \frac{f'(z)}{\varphi^n(z)} > 0 \quad (1)$$

Clearly $T_2 = K(1)$, the class of close-to-convex functions and it is easily seen [5] that $T_k \subset K(k/2)$ for $k \geq 2$

For $f \in K(1)$, Clunie and Pommerenke [1] showed that for $n \geq 2$, $n | a_n | < (2 + \sqrt{2})e M(n/(n+1))$, where $M(r) = \max_{\theta} |f(re^{i\theta})|$ and the author [7] showed that $L(r) < AM(r) \log 1/(1-r)$, where $L(r)$ denotes the length of the image of $\{z : |z| = r\}$ by $f(z)$ and where A is an absolute constant. The object of the

present paper is to extend these results to the class T_k . The question of whether the results remain valid in the wider class $K(\beta)$ for $\beta > 1$ remains open.

2. Results. THEOREM 1. *Let $f \in T_k$ ($k \geq 2$), with $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$. Then for $n \geq 2$,*

$$n |a_n| \leq 3k e M(n/(n+1)) \quad (2)$$

Proof. We modify the method of Clunie and Pommerenke [1]. From (1) write

$$z f'(z) = g(z)h(z), \quad (3)$$

so that $g \in R_k$, $h(0) = 1$ and $\Re h(z) > 0$ for $z \in D$.

Thus we can write $z f'(z) = 2g(z)\Re h(z) - g(z)\overline{h(z)}$. Now with $z = re^{i\theta}$,

$$\begin{aligned} na_n &= \frac{1}{2\pi r^n} \int_0^{2\pi} z f'(z) e^{-in\theta} d\theta \\ &= \frac{1}{\pi r^n} \int_0^{2\pi} g(z)\Re[h(z)]e^{-in\theta} d\theta - \frac{1}{2\pi r^n} \int_0^{2\pi} g(z)\overline{h(z)}e^{-in\theta} d\theta. \end{aligned}$$

Therefore

$$\begin{aligned} n |a_n| &\leq \frac{1}{\pi r^n} \int_0^{2\pi} |g(z)| \Re[h(z)] d\theta + \frac{1}{2\pi r^n} \left| \int_0^{2\pi} \overline{g(z)h(z)} e^{-in\theta} d\theta \right| \\ &= I_1(r) + I_2(r) \quad \text{say} \end{aligned}$$

Since $\Re h(z) > 0$ for $z \in D$, (3) gives

$$|g(z)| \Re[h(z)] = \Re[z f'(z) e^{-i \arg g(z)}].$$

Thus integrating by parts

$$I_1(r) = \frac{1}{\pi r^n} \Re \int_0^{2\pi} f(z) e^{-1 \arg g(z)} d_\theta(\arg g(z)) \leq \frac{k}{r^n} M(r),$$

since

$$\int_0^{2\pi} \left| \Re \frac{z g'(z)}{g(z)} \right| d\theta \leq k\pi \quad (4)$$

For $I_2(r)$, we have from (3)

$$I_2(r) = \frac{1}{2\pi r^{2n}} \left| \int_0^{2\pi} z^{n+1} f'(z) e^{-2i \arg g(z)} d\theta \right|. \quad (5)$$

Let $f_n(z) = \int_0^z t^n f'(t) dt$. Then integrating by parts gives

$$|f_n(z)| \leq 2r^n M(r). \quad (6)$$

Finally integrating by parts in (5) shows that

$$I_2(r) = \frac{1}{\pi r^{2n}} \left| \int_0^{2\pi} f_n(z) e^{-2i \arg g(z)} \Re \frac{z g'(z)}{g(z)} d\theta \right| \leq \frac{2k}{r^n} M(r)$$

on using (4) and (6).

Choosing $r = n/(n + 1)$ gives (2).

THEOREM 2. *Let $f \in T_k (k \geq 2)$. Then for $0 < r < 1$,*

$$L(r) \leq A(k)M(r) \log 1/(1 - r),$$

where $A(k)$ is a constant depending only upon k .

Proof. With $z = re^{i\theta}$, (3) gives

$$\begin{aligned} L(r) &= \int_0^r \int_0^{2\pi} |zf'(z)| d\theta \leq \int_0^r \int_0^{2\pi} |g'(\rho e^{i\theta})h(\rho e^{i\theta})| d\theta d\rho \\ &\quad + \int_0^r \int_0^{2\pi} |g(\rho e^{i\theta})h'(\rho e^{i\theta})| d\theta d\rho = J_1(r) + J_2(r) \quad \text{say.} \end{aligned}$$

Now $J_1(r) = \int_0^r \int_0^{2\pi} |f'(\rho e^{i\theta})H(\rho e^{i\theta})| d\theta d\rho$, where $H(z) = \frac{zg'(z)}{g(z)}$. Thus

$$\begin{aligned} J_1(r) &\leq \int_0^r \left(\int_0^{2\pi} |f'(\rho e^{i\theta})|^2 d\theta \right)^{\frac{1}{2}} \left(\int_0^{2\pi} |H(\rho e^{i\theta})|^2 d\theta \right)^{\frac{1}{2}} d\rho \\ &\leq 2\pi \int_0^r \left(1 + \sum_{n=2}^{\infty} n^2 |a_n|^2 \rho^{2n-2} \right)^{\frac{1}{2}} \left(\frac{1 + (k^2 - 1)\rho^2}{1 - \rho^2} \right)^{\frac{1}{2}} d\rho \end{aligned} \quad (7)$$

where we have used the Cauchy-Schwartz inequality, Parseval's equality and Lemma 2 in [5].

If $f \in K(\beta)$, $0 \leq \beta \leq 1$, then f is univalent in D [3]. However for $\beta > 1$, f need not be finitely valent [4]. Thus to estimate the first expression in (7) we proceed as follows.

With $\rho = n/(n + 1)$, (2) gives

$$\sum_{n=2}^{\infty} n^2 |a_n|^2 \rho^{2n-2} \leq 9k^2 e^2 M(\sqrt{\rho})^2 \sum_{n=2}^{\infty} \rho^{n-2}. \quad (8)$$

It follows immediately from the definition of T_k that the class T_k forms a subset of a linear-invariant family of order $k/2 + 1$. Using Lemma 2.6 of [6] we deduce that $M(\sqrt{\rho}) < 2^{k+2}M(\rho)/\sqrt{\rho}$. Thus from (7) and (8) we have $J_1(r) < A(k)M(r) \log 1/(1 - r)$.

To estimate $J_2(r)$ we note that since $\Re h(z) > 0$ for $z \in D$, $|h'(\rho e^{i\theta})| \leq 2\Re h(\rho e^{i\theta})/(1 - \rho^2)$. Thus

$$J_2(r) \leq 2 \int_0^r \int_0^{2\pi} \frac{|g(\rho e^{i\theta})| \Re h(\rho e^{i\theta})}{1 - \rho^2} d\theta d\rho \leq 2k\pi \int_0^r \frac{M(\rho)}{1 - \rho^2} d\rho$$

as in the proof of Theorem 1. Combining the estimates for $J_1(r)$ and $J_2(r)$ gives Theorem 2.

Remark. The proof of Theorem 2 shows that in fact

$$L(r) \leq A(k) \int_0^r \frac{M(\rho)}{1-\rho} d\rho.$$

Thus if $f \in T_k$ and $M(r) < 1/(1-r)^\alpha$, $\alpha > 0$, then $L(r) < A(k, \alpha)/(1-r)^\alpha$, where $A(k, \alpha)$ denotes a constant depending only upon k and α .

REFERENCES

- [1] J. Clunie and Ch. Pommerenke, *On the coefficients of close-to-convex univalent functions*, J. London Math. Soc. **41** (1966), 161-165.
- [2] P. L. Duren, *Univalent Functions*, Springer-Verlag, 1983.
- [3] A. W. Goodman, *On close-to-convex functions of higher order*, Ann. Univ. Sci. Budapest Eötvös Sect. Math **25** (1972), 17-30.
- [4] A. W. Goodman, *A note on the Nashiro-Warschawski theorem*, J. Anal. **25** (1972), 401-408.
- [5] K. I. Noor, *On a generalisation of close-convexity*, Internat. J. Math. Sci. **6** (1983), 327-334.
- [6] Ch. Pommerenke, *Linear-invariante analytischer Functionen I*, Math. Ann. **155** (1964), 108-154.
- [7] D. K. Thomas, *On starlike and close-convex univalent functions*, J. Lond. Math. Soc. **42** (1967), 427-435.

Department of Mathematics
and Computer Science
Swansea SA2 8PP,
Wales, U. K.

(Received 29 09 1988)