

**A NOTE ON THE HOLOMORPHIC INVARIANTS  
OF TIAN–ZHU \***

ZHIQIN LU

In this short note, we compute the holomorphic invariants defined by Tian and Zhu [4] on smooth hypersurfaces of  $CP^n$ . The holomorphic invariants, which generalize the famous Futaki invariants [1], are obstructions towards the existence of Kähler–Ricci solitons.

For a Kähler manifold with the first positive Chern class, the existence of the Kähler–Ricci soliton can be reduced to the existence of the solution of a nonlinear equation of Monge–Ampere type. In general, solving such an equation is highly nontrivial. Similarly to the Futaki invariants, the Tian–Zhu invariants give the obstruction *before* one needs to solve the equation. It is thus very important to compute it concretely. In this paper, in the case of hypersurfaces, we give an explicit formula.

Let  $M \subset CP^n$  be a smooth hypersurface defined by a homogeneous polynomial  $F = 0$  of degree  $d$ . Let  $v$  and  $X$  be two holomorphic vector fields on  $CP^n$ . For the sake of simplicity, we assume that

$$v = \sum_{i=0}^n v^i Z_i \frac{\partial}{\partial Z_i} \quad \text{and} \quad X = \sum_{i=0}^n X^i Z_i \frac{\partial}{\partial Z_i} ,$$

where  $[Z_0, \dots, Z_n]$  is the homogeneous coordinate of  $CP^n$ ,  $(v^0, \dots, v^n) \in C^{n+1}$ ,  $(X^0, \dots, X^n) \in C^{n+1}$ . We further assume that

$$(1) \quad \sum_{i=0}^n v^i = 0, \quad \sum_{i=0}^n X^i = 0 .$$

---

*Received:* December 18, 2001; *Revised:* March 13, 2002.

*AMS Subject Classification:* Primary 53A30; Secondary 32C16.

*Keywords and Phrases:* Kähler–Ricci soliton; Futaki invariants; Kähler–Einstein metric.

\* Research supported by NSF grant DMS 0196086.

If  $v$  and  $X$  are tangent vector fields of  $M$ , then there are complex numbers  $\lambda$  and  $\kappa$  such that

$$(2) \quad vF = \kappa F, \quad XF = \lambda F.$$

Let  $\omega$  be the Kähler form of the Fubini–Study metric of  $CP^n$ . Then  $(n-d+1)\omega$  restricts to a representative of the first Chern class  $c_1(M)$  of  $M$ . Thus there is a smooth function  $\xi$  on  $M$  such that

$$\text{Ric}((n-d+1)\omega|_M) - (n-d+1)\omega|_M = \partial\bar{\partial}\xi.$$

For fixed holomorphic vectors  $X$  and  $v$ , the holomorphic invariant defined by Tian–Zhu [4], in our context, is

$$(3) \quad F_X(v) = (n-d+1)^{n-1} \int_M v(\xi - (n-d+1)\theta_X) e^{(n-d+1)\theta_X} \omega^{n-1},$$

where  $\theta_X$  is defined as

$$(4) \quad \begin{cases} i(X)\omega = \frac{\sqrt{-1}}{2\pi} \bar{\partial}\theta_X, \\ \int_M e^{(n-d+1)\theta_X} \omega^{n-1} = d. \end{cases}$$

The main property of the Tian–Zhu invariants is the following (cf. [4]):

**Theorem 1.** *Let  $F_X(v)$  be the Tian–Zhu invariant. Then we have*

1. *If the Kähler–Ricci soliton exists, that is, we have*

$$\text{Ric}(\omega) - \omega = L_X\omega$$

*for some Kähler metric  $\omega$ . Then  $F_X(v) \equiv 0$ .*

2.  *$F_X(v)$  is independent of the choice of the Kähler metric  $\omega$  within the first Chern class.*

In this note, we give a “computable” expression of  $F_X(v)$ . Our main result is as follows:

**Theorem 2.** *Using the notations as above, defined the function*

$$(5) \quad \varphi(X) = \sum_{k=0}^{\infty} \frac{n!(n-d+1)^k}{(n+k)!} \sum_{\alpha_0+\dots+\alpha_n=k} X_0^{\alpha_0} \dots X_n^{\alpha_n},$$

where  $\alpha_0, \dots, \alpha_n \in \mathbb{Z}^{n+1}$  are nonnegative integers. Let

$$(6) \quad \sigma(X) = \left( -\frac{\lambda(n-d+1)}{n} + d \right) \varphi(X) + \frac{d}{n} \sum_{i=0}^n X^i \frac{\partial \varphi(X)}{\partial X^i} .$$

Then the invariants defined by Tian–Zhu can be explicitly expressed as

$$(7) \quad F_X(v) = -(n-d+1)^{n-1} d \left( \kappa + \sum_{i=0}^n v^i \frac{\partial \log \sigma(X)}{\partial X^i} \right) .$$

**Corollary 1.** *The Futaki invariant for the hypersurface  $M$  is*

$$F(v) = -(n-d+1)^{n-1} \frac{(n+1)(d-1)}{n} \kappa . \blacksquare$$

The rest of this note is devoted to the proof Theorem 2. We define

$$(8) \quad \tilde{\theta}_X = \frac{\lambda_0 |Z_0|^2 + \dots + \lambda_n |Z_n|^2}{|Z_0|^2 + \dots + |Z_n|^2} .$$

Then we have

$$(9) \quad i(X)\omega = \bar{\partial} \tilde{\theta}_X .$$

By comparing the above equation with (4), we have

$$(10) \quad \theta_X = \tilde{\theta}_X + c_X$$

for a constant  $c_X$ . First, we have the following lemma

**Lemma 1.**

$$\int_{CP^n} e^{(n-d+1)\tilde{\theta}_X} \omega^n = \varphi(X) ,$$

where  $\varphi(X)$  is defined in (5).

**Proof:** This follows from the expansion

$$e^{(n-d+1)\tilde{\theta}_X} = \sum_{k=0}^{\infty} \frac{(n-d+1)^k}{k!} \tilde{\theta}_X^k ,$$

and the elementary Calculus.  $\blacksquare$

**Lemma 2.** *Using the same notation as above, we have*

$$F_X(v) = (n - d + 1)^{n-1} \left( -\kappa d - \int_M (n - d + 1) \theta_v e^{(n-d+1)\theta_X} \omega^{n-1} \right).$$

**Proof:** By [3, Theorem 4.1], we have

$$\operatorname{div} v + v(\xi) + (n - d + 1)\theta_v = -\kappa,$$

where  $\theta_v$  is the function on  $CP^n$  defined by

$$\theta_v = \frac{v_0|Z_0|^2 + \cdots + v_n|Z_n|^2}{|Z_0|^2 + \cdots + |Z_n|^2},$$

and  $\kappa$  is defined in (2). Then (3) becomes

$$(11) \quad F_X(v) = (n - d + 1)^{n-1} \cdot \left( \int_M \left( -\kappa - \operatorname{div} v - (n-d+1)\theta_v - (n-d+1)v(\theta_X) \right) e^{(n-d+1)\theta_X} \omega^{n-1} \right).$$

We also have

$$(12) \quad \operatorname{div} (e^{(n-d+1)\theta_X} v) = e^{(n-d+1)\theta_X} \left( \operatorname{div} v + (n - d + 1) v(\theta_X) \right).$$

The lemma follows from (4), (11), (12) and the divergence theorem. ■

The following key lemma transfers the integration on  $M$  to the integrations on  $CP^n$ .

**Lemma 3.**

$$(13) \quad (n - d + 1) \int_M \theta_v e^{(n-d+1)\theta_X} \omega^{n-1} = d \sum_{i=0}^n v^i \frac{\partial \log \sigma}{\partial X^i},$$

where  $\sigma(X)$  is defined in (6).

**Proof:** Let

$$(14) \quad \eta = \log \frac{|F|^2}{\left( |Z_0|^2 + \cdots + |Z_n|^2 \right)^d}.$$

Then  $\eta$  is a smooth function on  $CP^n$  outside  $M$ . We have the following identity:

$$(15) \quad \begin{aligned} & \bar{\partial} \left( e^{(n-d+1)\theta_X} \partial \eta \wedge \omega^{n-1} \right) - \frac{n-d+1}{n} i(X) \left( e^{(n-d+1)\theta_X} \partial \eta \wedge \omega^n \right) = \\ & = -e^{(n-d+1)\theta_X} \partial \bar{\partial} \eta \wedge \omega^{n-1} - \frac{n-d+1}{n} e^{(n-d+1)\theta_X} (\lambda - d\tilde{\theta}_X) \omega^n. \end{aligned}$$

Since on  $CP^n$ , there are no  $(2n+1)$  forms, the left hand side of the above equation is the divergence of some vector field. Integrate the equation on both side and use the divergence theorem, we have

$$(16) \quad \int_{CP^n} e^{(n-d+1)\theta_X} \partial\bar{\partial}\eta \wedge \omega^{n-1} = -\frac{n-d+1}{n} \int_{CP^n} (\lambda - d\tilde{\theta}_X) e^{(n-d+1)\theta_X} \omega^n .$$

By [2, page 388], in the sense of currents, we have

$$(17) \quad \partial\bar{\partial}\eta = [M] - d\omega .$$

Thus from (16),

$$(18) \quad \begin{aligned} \int_M e^{(n-d+1)\theta_X} \omega^{n-1} &= \left( -\frac{\lambda(n-d+1)}{n} + d \right) \int_{CP^n} e^{(n-d+1)\theta_X} \omega^n \\ &+ \frac{d(n-d+1)}{n} \int_{CP^n} \tilde{\theta}_X e^{(n-d+1)\theta_X} \omega^n . \end{aligned}$$

From Lemma 1, we have

$$(19) \quad \sum_{i=0}^n X^i \frac{\partial\varphi(X)}{\partial X^i} = (n-d+1) \int_{CP^n} \tilde{\theta}_X e^{(n-d+1)\tilde{\theta}_X} \omega^n .$$

By (10), (18) and (19)

$$(20) \quad \int_M e^{(n-d+1)\theta_X} \omega^{n-1} = \sigma(X) e^{cX} .$$

From the above equation, we have

$$(21) \quad (n-d+1) \int_M \theta_v e^{(n-d+1)\theta_X} \omega^{n-1} = \sum_{i=0}^n v^i \frac{\partial\sigma(X)}{\partial X^i} e^{cX} .$$

On the other hand, from (20), we have

$$(22) \quad d = \sigma(X) e^{cX} ,$$

by (4). Lemma 3 follows from (21) and (22). ■

Theorem 2 follows from Lemma 2 and Lemma 3.

## REFERENCES

- [1] FUTAKI, A. – An obstruction to the existence problem of positive Kähler Einstein metrics, *Invent. Math.*, 73 (1983), 437–443.
- [2] GRIFFITHS, P. and HARRIS, J. – *Principle of Algebraic Geometry*, John Wiley & Sons, 1978.
- [3] LU, Z. – On the Futaki invariants of complete intersections, *Duke Math. J.*, 100(2) (1999), 359–372.
- [4] TIAN, G. and ZHU, X. – Uniqueness of Kähler–Ricci solitons, *Acta Math.*, 184 (2000), 271–305.

Zhiqin Lu,  
Department of Mathematics, University of California, Irvine,  
Irvine, CA 92697 – U.S.A.  
E-mail: [zlu@math.uci.edu](mailto:zlu@math.uci.edu)