

## FREE $Q$ -DISTRIBUTIVE LATTICE OVER AN $n$ -ELEMENT CHAIN

LUIZ MONTEIRO, MANUEL ABAD and MARTA ZANDER \*

**Abstract:** In this note we provide an explicit construction of  $FQ(\mathbf{n})$ , the free  $Q$ -distributive lattice over an  $n$ -element chain, different from those given by Cignoli [4] and Abad–Díaz Varela [1], and prove that  $FQ(\mathbf{n})$  can be endowed with a structure of a De Morgan algebra.

### 1 – Preliminaries

Quantifiers on distributive lattices were considered for the first time by Servi in [14], but it was Cignoli [3] who studied them as algebras, which he named  $Q$ -distributive lattices. In [4], Cignoli gave a construction of the free  $Q$ -distributive lattice over a set  $X$ . This construction generalizes that given by Halmos [6] for the free monadic Boolean algebra. The present paper is motivated by a result of Abad and Díaz Varela [1] on the free  $Q$ -distributive lattice,  $FQ(I)$ , over a finite ordered set  $I$ . They proved that  $FQ(\mathbf{n}) \cong \mathbf{2}^{[\mathbf{2}^{[2^{\times \mathbf{n}}]}]}$ , where  $\mathbf{n}$  is an  $n$ -element chain. We provide an easy construction for the diagram of  $\mathbf{2}^{[2^{\times \mathbf{n}}]}$ , and we prove some properties of  $FQ(\mathbf{n})$ .

We include in this section some notation, definitions and results on distributive lattices which are used in the paper.

An element  $p$  of a lattice  $R$  is said to be join-irreducible if  $p \neq 0$  and  $p = x \vee y$  implies  $p = x$  or  $p = y$ , for all  $x, y \in R$ . We denote the set of join-irreducible elements of  $R$  by  $\mathcal{J}(R)$ .

---

*Received:* May 9, 2002; *Revised:* March 12, 2003.

*AMS Subject Classification:* 08B20 06D99, 06D30.

*Keywords:* ordered sets;  $Q$ -distributive lattices; free algebras; De Morgan algebras.

\*The support of CONICET is gratefully acknowledged.

The following result was proved by Antonio Monteiro (see [8], [12]) in his lectures held at the Universidad Nacional del Sur.

**Lemma 1.1.** *If  $R$  is a finite distributive lattice and  $p \in R \setminus \{0\}$ , then  $p \in \mathcal{J}(R)$  if and only if there is a greatest element in the set  $I(p) = \{x \in R : x < p\}$ . ■*

For  $n \in \mathbb{N}$ , we write  $\mathbf{n}$  to denote the chain obtained by giving the  $n$ -element set  $\{1, 2, \dots, n\}$  the order in which  $1 < 2 < \dots < n$ .

Given an ordered set  $X$ , we denote the dual of  $X$  by  $X^*$  and the bounded distributive lattice of all order-preserving functions from  $X$  into  $\mathbf{2}$  by  $\mathbf{2}^{[X]}$  (see [7]).

A subset  $Y$  of  $X$  is said to be a *down-set*, or *order ideal* if, whenever  $b \in Y$ ,  $a \in X$  and  $a \leq b$ , we have  $a \in Y$ . Given  $x \in X$ , it is clear that the set  $(x) = \{y \in X : y \leq x\}$  is an order ideal; it is known as the principal order ideal generated by  $x$ . The notions of an *upper-set*, or *order filter* and of  $[x)$  are dually defined.

If  $X$  is a finite ordered set, the lattice  $\mathbf{2}^{[X]}$  is anti-isomorphic to the distributive lattice  $\mathcal{O}(X)$  of all down-sets of  $X$ , and  $\mathcal{J}(\mathcal{O}(X)) = \{(x) : x \in X\}$ .

**Lemma 1.2** (G. Birkhoff [2]). *For every finite ordered set  $X$ ,  $\mathcal{J}(\mathbf{2}^{[X]}) \cong X^*$ , and conversely, for every finite distributive lattice  $R$ ,  $R \cong \mathbf{2}^{[\mathcal{J}(R)^*]}$ . ■*

We use the following notations: the cardinality of a finite set  $X$  is denoted by  $N[X]$ , and the  $(0, 1)$ -sublattice generated by a subset  $X$  of a lattice  $R$  is denoted by  $SL(X)$ .  $FD(G)$  denotes the free bounded distributive lattice generated by  $G$ . An ordered set  $X$  is called *self-dual* if  $X$  and  $X^*$  are isomorphic ordered sets.

The following result can be found in B. Jónsson [7].

**Lemma 1.3.**  $FD(\mathbf{2} \times \mathbf{n}) \cong \mathbf{2}^{[\mathbf{2}^{[\mathbf{2} \times \mathbf{n}]}]}$ . ■

## 2 – Free $Q$ -distributive lattice over an $n$ -element chain

The aim of this section is to give a construction of the free  $Q$ -distributive lattice  $FQ(\mathbf{n})$  over an  $n$ -element chain. A  $Q$ -distributive lattice is an algebra  $\langle A, \wedge, \vee, 0, 1, \nabla \rangle$  of type  $(2, 2, 0, 0, 1)$  such that  $\langle A, \wedge, \vee, 0, 1 \rangle$  is a bounded distributive lattice and  $\nabla$  is an existential quantifier, that is,  $\nabla 0 = 0$ ,  $x \leq \nabla x$ ,  $\nabla(x \wedge \nabla y) = \nabla x \wedge \nabla y$ ,  $\nabla(x \vee y) = \nabla x \vee \nabla y$ .

We introduce an ordered set  $V(n)$  isomorphic to  $\mathbf{2}^{[2 \times \mathbf{n}]}$ , so that

$$\mathcal{O}(V(n)) \cong \mathbf{2}^{[2 \times \mathbf{n}]} ,$$

and define an operator  $\nabla$  on  $\mathcal{O}(V(n))$ , in such a way that  $\langle \mathcal{O}(V(n)), \nabla \rangle$  is isomorphic to the free  $Q$ -distributive lattice over an  $n$ -element chain.

Let  $V(n) = \{(x, y) \in (\mathbf{n} + \mathbf{1}) \times (\mathbf{n} + \mathbf{1}) : x \leq y\}$ . It is clear that  $V(n)$  is a  $(0, 1)$ -sublattice of  $(\mathbf{n} + \mathbf{1}) \times (\mathbf{n} + \mathbf{1})$ . Observe that  $V(n)$  is self-dual and

$$N[V(n)] = \frac{(n + 1)(n + 2)}{2} .$$

Now we want to characterize the set  $\mathcal{J}(V(n))$ .

For every  $j$ ,  $2 \leq j \leq n + 1$ ,  $I((1, j)) = ((1, j - 1))$ . So, from Lemma 1.1, we have that  $(1, j) \in \mathcal{J}(V(n))$ . If  $i \geq 2$ , then  $I((i, i)) = ((i - 1, i))$ . So, again from Lemma 1.1,  $(i, i) \in \mathcal{J}(V(n))$ . Since  $(i, j) = (i, i) \vee (1, j)$ , it follows that

$$\mathcal{J}(V(n)) = \bigcup_{j=2}^{n+1} \{(1, j)\} \cup \bigcup_{i=2}^{n+1} \{(i, i)\} .$$

It is clear that the ordered sets  $\mathcal{J}(V(n))$  and  $\mathbf{2} \times \mathbf{n}$  are isomorphic, and consequently the distributive lattices  $V(n)$  and  $\mathbf{2}^{[2 \times \mathbf{n}]}$  are isomorphic. From Lemma 1.3,

$$FD(\mathbf{2} \times \mathbf{n}) \cong \mathbf{2}^{[V(n)]} .$$

Let us see now that  $N[\mathbf{2}^{[V(n)]}] = 2^{n+1}$ , for every  $n \in \mathbb{N}$ . In the proof we make use of the following result of L. Monteiro [13]:

**Lemma 2.1.** *If  $X$  is a finite ordered set and  $f$  is an element of  $X$  which is neither first nor last element of  $X$ , then*

$$N[\mathbf{2}^{[X]}] = N[\mathbf{2}^{[X \setminus \{f\}]}] + N[\mathbf{2}^{[X \setminus \{f\}]}] . \blacksquare$$

To prove that  $N[\mathbf{2}^{[V(n)]}] = 2^{n+1}$  we proceed by induction. For  $n = 1$ ,  $N[\mathbf{2}^{[V(1)]}] = N[\mathbf{2}^{[V(1) \setminus \{(1,2)\}]}] + N[\mathbf{2}^{[V(1) \setminus \{(1,2)\}]}] = 2 + 2 = 2^{1+1}$ . Suppose that  $N[\mathbf{2}^{[V(n-1)]}] = 2^n$ , for  $n \geq 2$ . Then

$$N[\mathbf{2}^{[V(n)]}] = N[\mathbf{2}^{[V(n) \setminus \{(1, n+1)\}]}] + N[\mathbf{2}^{[V(n) \setminus \{(1, n+1)\}]}] .$$

But  $V(n) \setminus \{(1, n + 1)\}$  and  $V(n) \setminus \{(1, n + 1)\}$  are ordered sets isomorphic to  $V(n - 1)$ . So  $N[\mathbf{2}^{[V(n)]}] = 2 \cdot N[\mathbf{2}^{[V(n-1)]}] = 2 \cdot 2^n = 2^{n+1}$ .

Let  $m: V(n) \rightarrow V(n)$  be the map defined by  $m(i, j) = (i, n + 1)$ .  $m$  is a lattice homomorphism that satisfies  $x \leq m(x)$  and  $m(m(x)) = m(x)$ .

Now we define the operator  $\nabla$  on  $\mathcal{O}(V(n))$ . For  $X \in \mathcal{O}(V(n))$ , let

$$\nabla X = \begin{cases} \emptyset, & \text{if } X = \emptyset \\ \bigcup_{x \in X} (m(x)], & \text{if } X \in \mathcal{O}(V(n)) \setminus \{\emptyset\}. \end{cases}$$

Let us prove that  $\langle \mathcal{O}(V(n)), \nabla \rangle$  is a  $Q$ -distributive lattice.

By the definition,  $\nabla \emptyset = \emptyset$ .

Since  $x \leq m(x)$ , it follows that  $\nabla X = \bigcup_{x \in X} (m(x)] \supseteq \bigcup_{x \in X} (x] = X$ .

$$\nabla(X \cup Y) = \bigcup_{z \in X \cup Y} (m(z)] = \bigcup_{z \in X} (m(z)] \cup \bigcup_{z \in Y} (m(z)] = \nabla X \cup \nabla Y.$$

Finally, let us see that  $\nabla(X \cap \nabla Y) = \nabla X \cap \nabla Y$ . If  $t \in \nabla(X \cap \nabla Y)$  then  $t \leq m(w)$ ,  $w \in X \cap \nabla Y$ . As  $w \in X$ , then  $t \in \nabla X$ . Since  $w \in \nabla Y$ , we have that  $w \leq m(y)$ , for some  $y \in Y$ , and consequently,  $t \leq m(w) \leq m(m(y)) = m(y)$ . Thus  $t \in \nabla Y$ . Conversely, if  $t \in \nabla X \cap \nabla Y$ , then  $t \leq m(x)$ ,  $x \in X$  and  $t \leq m(y)$ ,  $y \in Y$ . Since  $m$  is a homomorphism,  $t \leq m(x \wedge y)$ , where  $x \wedge y \in X \cap Y \subseteq X \cap \nabla Y$ , that is,  $t \in \nabla(X \cap \nabla Y)$ .

**Theorem 2.1.**  $\langle \mathcal{O}(V(n)), \nabla \rangle$  is the free  $Q$ -distributive lattice over  $\mathbf{n}$ .

**Proof:** First we prove that  $\langle \mathcal{O}(V(n)), \nabla \rangle$  is generated by the chain

$$G = \{g_i = ((i, i]): 1 \leq i \leq n\}.$$

Observe that

$$\nabla g_j = \nabla((j, j]) = \bigcup_{x \in ((j, j])} (m(x)] = (m(j, j)] = ((j, n + 1)], \quad 1 \leq j \leq n.$$

So

$$G \cup \nabla G = \{((i, i]): 1 \leq i \leq n\} \cup \{((i, n + 1]): 1 \leq i \leq n\}.$$

If  $(i, j) \in V(n)$ , then  $(i, j) = (j, j) \wedge (i, n + 1)$ . Hence, if  $j \neq n + 1$ ,

$$((i, j]) = ((j, j]) \cap ((i, n + 1]) = g_j \cap \nabla g_i.$$

That is, every principal down-set of  $V(n)$  different from  $V(n)$  is a meet of elements in  $G \cup \nabla G$ . Since every non-empty element of  $\mathcal{O}(V(n))$  is a union of principal

down-sets of  $V(n)$ , we have that every element of  $\mathcal{O}(V(n))$  different from  $\emptyset$  and  $V(n)$  is a union of meets of elements of  $G \cup \nabla G$ , that is,

$$SL(G \cup \nabla G) = \mathcal{O}(V(n)) .$$

So it is clear that  $\mathcal{O}(V(n))$  is generated by  $G$  as a  $Q$ -distributive lattice.

In order to prove that  $\mathcal{O}(V(n))$  is the free  $Q$ -distributive lattice over  $\mathbf{n}$ , let  $f: \mathbf{n} \rightarrow \mathcal{O}(V(n))$  be the mapping defined by  $f(i) = g_i$ ,  $1 \leq i \leq n$ .  $f$  can be extended to a  $Q$ -homomorphism  $\bar{f}: FQ(\mathbf{n}) \rightarrow \mathcal{O}(V(n))$ . Now,  $\bar{f}(FQ(\mathbf{n}))$  is a  $Q$ -sublattice of  $\mathcal{O}(V(n))$  such that  $G \subseteq \bar{f}(FQ(\mathbf{n}))$  and since  $\mathcal{O}(V(n))$  is the  $Q$ -distributive lattice generated by  $G$ , we have  $\bar{f}(FQ(\mathbf{n})) = \mathcal{O}(V(n))$ . So  $\bar{f}$  is a  $Q$ -epimorphism. Since the sets  $FQ(\mathbf{n})$  and  $\mathcal{O}(V(n))$  are finite and they have the same cardinality,  $\bar{f}$  is also injective. Consequently,  $\bar{f}$  is an isomorphism. ■

### 3 – A structure of De Morgan algebra on $FQ(\mathbf{n})$

The aim of this section is to define a De Morgan negation on  $\mathcal{O}(V(n))$  in order to establish a relationship between  $\mathcal{O}(V(n))$  and  $\mathcal{O}(V(n-1))$ .

Since  $V(n)$  is self-dual, it is possible to define a De Morgan operation on  $\mathcal{O}(V(n))$ . Indeed, let  $\varphi: V(n) \rightarrow V(n)$  be defined by

$$\varphi(x, y) = (n + 2 - y, n + 2 - x) .$$

It is clear that  $\varphi$  is an anti-isomorphism of period 2. So we can define on  $\mathcal{O}(V(n))$  a De Morgan operation  $\sim$  associated to  $\varphi$  in the usual way [9, 10, 11]: if  $Y \in \mathcal{O}(V(n))$  then

$$\sim Y = \bigcup \{(\varphi(p)) : p \in V(n), p \notin Y\} .$$

Let us see that

$$(3.1) \quad \sim g_j = \nabla g_{n+1-j}, \quad \text{for } 1 \leq j \leq n .$$

First observe that  $\varphi((p)) = [\varphi(p)]$  and  $\varphi([p]) = (\varphi(p))$  for every  $p \in V(n)$ . Thus

$$\begin{aligned} \sim g_j &= \sim ((j, j)) \\ &= \bigcup \{(\varphi(p)) : p \in V(n), p \notin ((j, j))\} \\ &= \bigcup \{\varphi([p]) : p \in V(n), p \notin ((j, j))\} \\ &= \varphi\left(\bigcup \{[p] : p \in V(n), p \notin ((j, j))\}\right) \\ &= \varphi([(1, j+1)]) = (\varphi(1, j+1)) = ((n+1-j, n+1)) = \nabla g_{n+1-j} . \end{aligned}$$

**Theorem 3.1.**  $\langle \mathcal{O}(V(n)), \sim \rangle$  is a Kleene algebra.

**Proof:** If  $A$  is a De Morgan algebra and  $X \subseteq A$ , let  $SM(X)$  denote the De Morgan subalgebra of  $A$  generated by  $X$ . It is known that  $SM(X) = SL(X \cup \sim X)$ . So, as  $G = \{g_1, g_2, \dots, g_n\} \subseteq \mathcal{O}(V(n))$ , by (3.1) we have that  $SM(G) = SL(G \cup \nabla G) = \mathcal{O}(V(n))$ . Hence the De Morgan algebra  $\mathcal{O}(V(n))$  is generated by an  $n$ -element chain, and consequently it is a homomorphic image of the free De Morgan algebra  $\mathcal{M}$  over an  $n$ -element chain.  $\mathcal{M}$  is a Kleene algebra [5], so  $\langle \mathcal{O}(V(n)), \sim \rangle$  is a Kleene algebra. ■

Now we are going to prove that:

**Theorem 3.2.** For  $n \geq 2$ , the down-set  $(g_n]$  and the upper-set  $[\nabla g_1)$  of  $\mathcal{O}(V(n))$  are ordered sets isomorphic to  $\mathcal{O}(V(n-1))$ .

**Proof:** Since  $\sim$  is an order anti-isomorphism from  $\mathcal{O}(V(n))$  on  $\mathcal{O}(V(n))$ , and

$$X \in (g_n] \quad \text{iff} \quad X \subseteq g_n \quad \text{iff} \quad \nabla g_1 = \sim g_n \subseteq \sim X,$$

then  $\sim$  induces an anti-isomorphism from  $(g_n]$  on  $[\nabla g_1)$ , that is,  $(g_n] \cong [\nabla g_1)^*$ .

Now observe that  $\mathcal{J}((g_n]) = \mathcal{J}(\mathcal{O}(V(n))) \cap (g_n]$ , and this is clearly isomorphic to  $V(n-1)$ . Then  $(g_n] \cong \mathcal{O}(V(n-1))$ . In particular,  $(g_n] \cong (g_n]^*$ , thus


$$[\nabla g_1) \cong \mathcal{O}(V(n-1)) \cong (g_n].$$

Observe that  $\mathcal{O}(V(n))$  is the disjoint union of  $[\nabla g_1)$  and  $(g_n]$ . Indeed, if  $X \not\subseteq g_n = ((n, n])$ , there exists  $(x_1, x_2) \in X$  such that  $(x_1, x_2) \notin ((n, n])$ , that is,  $(x_1, x_2) \not\subseteq (n, n)$ . Then  $(1, n+1) \leq (x_1, x_2)$ , and consequently,  $\nabla g_1 = ((1, n+1]) \subseteq X$ . So  $X \in [\nabla g_1)$ . If  $X \in [\nabla g_1) \cap (g_n]$  we have that  $\nabla g_1 = ((1, n+1]) \subseteq X$ . In particular,  $(1, n+1) \in X$  and  $X \subseteq g_n = ((n, n])$ . Thus  $(1, n+1) \in ((n, n])$ , a contradiction. ■


In particular,  $N[\mathcal{O}(V(n))] = 2 \cdot N[\mathcal{O}(V(n-1))]$ . This fact can be used to get another proof of the result  $N[\mathcal{O}(V(n))] = 2^{n+1}$ .

This situation is illustrated in the following figure.

$\mathcal{O}(V(3))$



$\mathcal{O}(V(4))$



## REFERENCES

- [1] ABAD, M. and DÍAZ VARELA, J.P. – Free  $Q$ -distributive lattices from meet semi-lattices, *Discrete Math.*, 224 (2000), 1–14.
- [2] BIRKHOFF, G. – *Lattice Theory*, AMS Colloquium Publ. 25, 3<sup>rd</sup> edn., Amer. Math. Soc. Providence, RI, 1967.
- [3] CIGNOLI, R. – Quantifiers on distributive lattices, *Discrete Math.*, 96 (1991), 183–197.
- [4] CIGNOLI, R. – Free  $Q$ -distributive lattices, *Studia Logica*, 56 (1996), 23–29.
- [5] FIGALLO, A. and MONTEIRO, L. – Système déterminant de l’algèbre de Morgan libre sur un ensemble ordonné fini, *Port. Math.*, 40(2) (1981), 129–135.
- [6] HALMOS, P. – Algebraic Logic I: monadic algebras, *Comp. Math.*, 12 (1955), 217–249. Reproduced in *Algebraic Logic*, Chelsea Pub. Co. New York, 1962.
- [7] JÓNSSON, B. – *Arithmetic of ordered sets*, in “Ordered Sets” (I. Rival, Ed.), D. Reidel Publishing Company, Dordrecht, Boston, pp. 3–41, 1982.
- [8] MONTEIRO, A. – *Notas del curso Algebra de la Lógica I*, Instituto de Matemática, Universidad Nacional del Sur, 1959.
- [9] MONTEIRO, A. – Matrices de Morgan caractéristiques pour le calcul propositionnel classique, *Anais da Academia Brasileira de Ciências*, 1–7 (1960), *Notas de Lógica Matemática*, 6–7, INMABB-CONICET-UNS, Bahía Blanca (1974), 39 pag.
- [10] MONTEIRO, A. – *Notas del curso Algebras de De Morgan*, Instituto de Matemática, Universidad Nacional del Sur, 1966.
- [11] MONTEIRO, A. – Sur les algèbres de Heyting symétriques, *Port. Math.*, 39(1)–(4) (1980), 1–237.
- [12] MONTEIRO, L. – *Algebras de Boole*, Informe Técnico Interno, 66 (2000), 197 pag., INMABB-CONICET-Universidad Nacional del Sur.
- [13] MONTEIRO, L. – Free bounded distributive lattices on  $n$  generators, *Multi. Val. Logic*, 6(1)–(2) (2001), 175–192.
- [14] SERVI, M. – Un’assiomatizzazione dei reticoli esistenziali, *Boll. Un. Mat. Ital. A*, 16(5) (1979), 298–301.

Luiz Monteiro, Manuel Abad and Marta Zander,  
INMABB-CONICET and Departamento de Matemática,  
Universidad Nacional del Sur – ARGENTINA

E-mail: [luizmont@criba.edu.ar](mailto:luizmont@criba.edu.ar)  
[imabad@criba.edu.ar](mailto:imabad@criba.edu.ar)  
[mzander@criba.edu.ar](mailto:mzander@criba.edu.ar)