

## Leafwise smoothing laminations

DANNY CALEGARI

**Abstract** We show that every topological surface lamination  $\Lambda$  of a 3-manifold  $M$  is isotopic to one with smoothly immersed leaves. This carries out a project proposed by Gabai in [2]. Consequently any such lamination admits the structure of a *Riemann surface lamination*, and therefore useful structure theorems of Candel [1] and Ghys [3] apply.

**AMS Classification** 57M50

**Keywords** Lamination, foliation, leafwise smooth, 3-manifold

### 1 Basic notions

**Definition 1.1** A *lamination* is a topological space which can be covered by open charts  $U_i$  with a local product structure  $\phi_i : U_i \rightarrow \mathbb{R}^n \times X$  in such a way that the manifold-like factor is preserved in the overlaps. That is, for  $U_i \cap U_j$  nonempty,

$$\phi_j \circ \phi_i^{-1} : \mathbb{R}^n \times X \rightarrow \mathbb{R}^n \times X$$

is of the form

$$\phi_j \circ \phi_i^{-1}(t, x) = (f(t, x), g(x))$$

The maximal continuations of the local manifold slices  $\mathbb{R}^n \times \text{point}$  are the *leaves* of the lamination. A *surface lamination* is a lamination locally modeled on  $\mathbb{R}^2 \times X$ . We usually assume that  $X$  is locally compact.

**Definition 1.2** A lamination is *leafwise  $C^n$*  for  $n \geq 2$  if the leafwise transition functions  $f(t, x)$  can be chosen in such a way that the mixed partial derivatives in  $t$  of orders less than or equal to  $n$  exist for each  $x$ , and vary continuously as functions of  $x$ .

A *leafwise  $C^n$  structure* on a lamination  $\Lambda$  induces on each leaf  $\lambda$  of  $\Lambda$  a  $C^n$  manifold structure, in the usual sense.

**Definition 1.3** An embedding of a leafwise  $C^n$  lamination  $i : \Lambda \rightarrow M$  into a manifold  $M$  is an  $C^n$  immersion if, for some  $C^n$  structure on  $M$ , for each leaf  $\lambda$  of  $\Lambda$  the embedding  $\lambda \rightarrow M$  is  $C^n$ .

Note that if  $i : \Lambda \rightarrow M$  is an embedding with the property that the image of each leaf  $i(\lambda)$  is locally a  $C^n$  submanifold, and these local submanifolds vary continuously in the  $C^n$  topology, then there is a unique leafwise  $C^n$  structure on  $\Lambda$  for which  $i$  is a  $C^n$  immersion.

A foliation of a manifold is an example of a lamination. For a foliation to be leafwise  $C^n$  is *a priori* weaker than to ask for it to be  $C^n$  immersed.

**Example 1.4** Let  $M$  be a manifold which is not stably smoothable, and  $N$  a compact smooth manifold. Then  $M \times N$  has the structure of a leafwise smooth foliation (by parallel copies of  $N$ ), but there is no smooth structure on  $M \times N$  for which the embedding of the foliation is a smooth immersion, since there is no smooth structure on  $M \times N$  at all.

**Remark 1.5** For readers unfamiliar with the notion, the “tangent bundle” of a topological manifold (i.e. a regular neighborhood of the diagonal in  $M \times M$ ) is stably (in the sense of  $K$ -theory) classified by a homotopy class of maps  $f : M \rightarrow BTOP$  for a certain topological space  $BTOP$ . There is a fibration  $p : BO \rightarrow BTOP$ , and the problem of lifting  $f$  to  $\hat{f} : M \rightarrow BO$  such that  $p\hat{f} = f$  represents an obstruction to finding a smooth structure on  $M$ . For  $N$  smooth as above, the composition

$$M \rightarrow M \times \text{point} \subset M \times N \rightarrow BTOP$$

is homotopic to  $f$ , and therefore no lifting of the structure exists on  $M \times N$  if none existed on  $M$ . For a reference, see [4], or the very readable [6].

With notation as above, the tangential quality of  $\mathcal{F}$  is controlled by the quality of  $f(\cdot, x)$  for each fixed  $x$ , for  $f$  the first component of a transition function. For sufficiently large  $k$  and  $n - k$  questions of ambiently smoothing *foliated manifolds* come down to obstruction theory and classical surgery theory, as for example in [4]. But in low dimensions, the situation is more elementary and more hands-on.

## 2 Some 3-manifold topology

Let  $M$  be a topological 3-manifold. It is a classical theorem of Moise (see [5]) that  $M$  admits a PL or smooth structure, unique up to conjugacy.

**Lemma 2.1** *Let  $\Sigma$  be a topological surface. Let  $S_j^1$  be a countable collection of circles, and let  $\Phi : \coprod_j S_j^1 \times I \rightarrow \Sigma$  be a map with the following properties:*

- (1) *For each  $t \in I$ ,  $\Phi(\cdot, t) : S_j^1 \rightarrow \Sigma$  is an embedding.*
- (2) *For each  $t \in I$  and each pair  $j, k$  the intersection*

$$\Phi(S_j^1, t) \cap \Phi(S_k^1, t)$$

*is finite, and its cardinality is constant as a function of  $t$  away from finitely many values.*

- (3) *For every compact subset  $K \subset \Sigma$  the set of  $j$  for which  $\Phi(S_j^1, t) \cap K$  is nonempty for some  $t$  is finite.*

*Then there is a PL (resp. smooth) structure on  $\Sigma \times I$  such that the graph of each map  $\Gamma_j(\Phi) : S_j^1 \times I \rightarrow \Sigma \times I$  is PL (resp. smooth).*

Here the *graph*  $\Gamma_j(\Phi)$  of  $\Phi$  is the function  $\Gamma_j(\Phi) : S_j^1 \times I \rightarrow \Sigma \times I$  defined by

$$\Gamma_j(\Phi)(\theta, t) = (\Phi(\theta), t)$$

**Proof** The conditions imply that the image of  $\coprod_j S_j^1$  in  $\Sigma$  for a fixed  $t$  is topologically a locally finite graph. Such a structure in a 2 manifold is locally flat, and the combinatorics of any finite subgraph is locally constant away from isolated values of  $t$ . It is therefore straightforward to construct a PL (resp. smooth) structure on a collar neighborhood of the image of  $\coprod_j S_j^1 \times I$  in  $\Sigma \times I$ . This can be extended canonically to a PL (resp. smooth) structure on  $\Sigma \times I$ , by the relative version of Moise’s theorem (see [5]). □

**Lemma 2.2** *Let  $\Phi : S_j^1 \times I \rightarrow \Sigma$  satisfy the conditions of lemma 2.1. Let  $\Psi_0 : S^1 \rightarrow \Sigma$  and  $\Psi_1 : S^1 \rightarrow \Sigma$  be homotopic embeddings such that  $\Psi_0(S^1)$  intersects finitely many circles in  $\Phi(\cdot, 0)$  in finitely many points, and similarly for  $\Psi_1(S^1)$ . Then there is a map  $\Psi : S^1 \times I \rightarrow \Sigma$  which is a homotopy between  $\Psi_0$  and  $\Psi_1$  so that*

$$\Phi \cup \Psi : \left( \coprod_i S_i^1 \coprod S^1 \right) \times I \rightarrow \Sigma$$

*satisfies the conditions of lemma 2.1.*

**Proof** Since the combinatorics of the image of  $\Phi$  is locally finite, and since the image of  $\Psi$  is bounded, it suffices to treat the case when  $\Phi$  is constant as a function of  $t$ .

Choose a PL structure on  $\Sigma$  for which the image of  $\Phi(\cdot, 0)$  and  $\Psi_0$  are polygonal. Then produce a polygonal homotopy from  $\Psi_0$  (with respect to this polygonal structure) to a new polygonal  $\Psi'_0$  such that  $\Psi'_0(S^1)$  and  $\Psi_1(S^1)$  intersect the image of  $\Phi(\cdot, t)$  in a finite set of points in the same combinatorial configuration. Then  $\Psi'_0$  is isotopic to  $\Psi_1$  rel. its intersection with the image of  $\Phi(\cdot, t)$ .  $\square$

### 3 Surface laminations of 3-manifolds

**Definition 3.1** Let  $\mathcal{F}$  be a codimension one foliation of a 3-manifold  $M$ . A *snake* in  $M$  is an embedding  $\phi : D^2 \times I \rightarrow M$  where  $D^2$  denotes the open unit disk, and  $I$  the open unit interval, which extends to an embedding of the closure of  $D^2 \times I$ , in such a way that each horizontal disk gets mapped into a leaf  $\lambda$  of  $\mathcal{F}$ . That is,  $\phi : D^2 \times t \rightarrow \lambda$ .

The terminology suggests that we are typically interested in snakes which are reasonably small and thin in the leafwise direction, and possibly large in the transverse direction.

A collection of snakes in a foliated manifold intersect a leaf  $\lambda$  of  $\mathcal{F}$  in a locally finite collection of open disks. For a snake  $S$ , let  $\partial_v \overline{S}$  denote the “vertical boundary” of the closed ball  $\overline{S}$ ; this is topologically an embedded closed cylinder transverse to  $\mathcal{F}$ , intersecting each leaf in an inessential circle.

We say that an open cover of  $M$  by finitely many snakes  $S_i$  is *combinatorially tame* if the embeddings  $\partial_v \overline{S}_i \rightarrow M$  are locally of the form described in lemma 2.1.

Note that the induced pattern on each leaf  $\lambda$  of  $\mathcal{F}$  of the circles  $\partial_v \overline{S}_i \cap \lambda$  is topologically conjugate to the transverse intersection of a locally finite collection of polygons.

**Lemma 3.2** *A codimension one foliation  $\mathcal{F}$  of a closed 3-manifold  $M$  admits a combinatorially tame open cover by finitely many snakes.*

**Proof** Since  $M$  is compact, any cover by snakes contains a finite subcover; any such cover induces a locally finite cover of each leaf. We prove the lemma by induction.

Let  $S_i$  be a collection of snakes in  $M$  which is combinatorially tame. Let  $C_i = \partial_v \overline{S}_i$  be their vertical boundaries, and let  $S$  be another snake with vertical

boundary  $C$ . We will show that there is a snake  $S'$  containing  $S$  such that the collection  $\{S_i\} \cup \{S'\}$  is combinatorially tame.

Let  $\lambda_t$  for  $t \in I$  parameterize the foliation of  $\overline{S}$ . Let  $E_i(t)$  denote the pattern of circles  $C_i \cap \lambda_t$  in a neighborhood of  $E(t) = C \cap \lambda_t$ . By hypothesis, the  $C_i$  can be thought of as polygons with respect to a PL structure on  $\lambda_t$ . Then  $E(t)$  can be *straightened* to a polygon  $E(t)'$  in general position with respect to the  $E_i(t)$  in a small neighborhood, where the interior of the region in  $\lambda_t$  bounded by  $E(t)'$  contains  $E(t)$ . If  $\lambda_t$  does not intersect the horizontal boundary of any  $\overline{S}_i$ , then the combinatorial pattern of intersections of the  $E_i(t)$  is locally generic — i.e. the pattern might change, but it changes by the graph of a generic PL isotopy, by lemma 2.1.

It follows that we can extend the straightening of  $E(t)$  to  $E(t)'$  for some collar neighborhood of  $t = 0$ . In general, a straightening of  $E(t)$  to  $E(t)'$  can be extended in the positive direction until a  $t_0$  which contains some lower horizontal boundary of an  $\overline{S}_i$ . The straightening can be extended past an upper horizontal boundary of an  $\overline{S}_i$  without any problems, since the combinatorial pattern of intersections becomes simpler: circles disappear.

The straightening of  $E(t)$  over all  $t$  can be done by *welding* straightenings centered at the finitely many values of  $t$  which contain horizontal boundary of some  $\overline{S}_i$ . Call these critical values  $t_j$ . So we can produce a finite collection of straightenings  $E(t) \rightarrow E(t)'_j$  each valid on the open interval  $t \in (t_{j-1}, t_{j+1})$ . To weld these straightenings together at intermediate values  $s_j$  where  $t_j < s_j < t_{j+1}$ , we insert a PL isotopy from  $E(s_j)'_j$  to  $E(s_j)'_{j+1}$  in a little collar neighborhood of  $s_j$ , by appealing to lemma 2.2. So these welded straightenings give a straightening of  $E(t)$  for all  $t \in I$ , and they bound a snake  $S'$  with the requisite properties.

To prove the lemma, cover  $M$  with finitely many snakes  $S_i$ , and apply the induction step to straighten  $S_j$  while fixing  $S_k$  with  $k < j$ . Since snakes can be straightened by an arbitrarily small (in the  $C^0$  topology) homotopy, the union of straightened snakes can also be made to cover  $M$ , and we are done.  $\square$

**Lemma 3.3** *Let  $M$  be a 3-manifold, and  $\mathcal{F}$  a foliation of  $M$  by surfaces. Then  $\mathcal{F}$  is isotopic to a foliation such that all leaves are PL or smoothly immersed, and the images of leaves vary locally continuously in the  $C^\infty$  topology.*

**Proof** If  $S_i$  is a combinatorially tame cover of  $\mathcal{F}$  by snakes, the image of the union  $\cup_i \partial \overline{S}_i$  can be taken to be a PL or smooth 2 complex  $\Sigma$  in  $M$ , whose complementary regions are polyhedral 3 manifolds. Each complementary region

is foliated as a product by  $\mathcal{F}$ . We can straighten  $\mathcal{F}$  cell-wise inductively on its intersection with the skeleta of  $\Sigma$ . First, we keep  $\mathcal{F} \cap \Sigma^1$  constant. Then the foliation of  $\mathcal{F} \cap (\Sigma^2 \setminus \Sigma^1)$  by lines can be straightened to be PL or smooth, and this straightened foliation extended in a PL or smooth manner over the product complementary regions in  $M - \Sigma$ .  $\square$

**Theorem 3.4** *Let  $\Lambda$  be a surface lamination in a 3-manifold  $M$ . Then  $\Lambda$  is isotopic to a lamination such that all leaves are PL or smoothly immersed, and the images of leaves vary locally continuously in the  $C^\infty$  topology.*

**Proof** By the definition of a lamination, there is an open cover of  $\Lambda$  by balls  $B_i$  such that  $\Lambda \cap B_i$  is a product lamination, which can be extended to a product foliation. It is straightforward to produce an open submanifold  $N$  with  $\Lambda \subset N \subset M$  such that  $N$  can be foliated by a foliation  $\mathcal{F}$  which contains  $\Lambda$  as a sublamination. Then the open manifold  $N$  can be given a PL or smooth structure in which  $\mathcal{F}$ , and hence  $\Lambda$ , is PL or smoothly immersed, by lemma 3.3. This PL or smooth structure can be extended compatibly over  $M - N$  by Moise's theorem.  $\square$

**Corollary 3.5** *Let  $\Lambda$  be a surface lamination in a 3-manifold  $M$ . Then  $\Lambda$  admits a leafwise PL or smooth structure.*

In particular, such a lamination admits the structure of a Riemannian surface lamination. In Gabai's problem list [2], he lists theorem 3.4 as a "project". The corollary allows us to apply the technology of complex analysis and algebraic geometry to such laminations; in particular, the following theorems of Candel and Ghys from [1] and [3] apply:

**Theorem 3.6** (Candel) *Let  $\mathcal{F}$  be an essential Riemann surface lamination of an atoroidal 3-manifold. Then there exists a continuously varying path metric on  $\mathcal{F}$  for which the leaves of  $\mathcal{F}$  are locally isometric to  $\mathbb{H}^2$ .*

**Theorem 3.7** (Ghys) *Let  $\mathcal{F}$  be a taut foliation of a 3-manifold  $M$  with Riemann surface leaves. Then there is an embedding  $e : M \rightarrow \mathbb{C}\mathbb{P}^n$  for some  $n$  which is leafwise holomorphic. That is,  $e : \lambda \rightarrow \mathbb{C}\mathbb{P}^n$  is holomorphic for each leaf  $\lambda$ .*

## References

- [1] A. Candel, *Uniformization of surface laminations*, Ann. Sci. École Norm. Sup. **26** (4) (1993) pp. 489–516
- [2] D. Gabai, *Problems in foliations and laminations*, in Geometric Topology (Athens, GA, 1993) AMS/IP Stud. Adv. Math. 2.2 pp. 1–33
- [3] E. Ghys, in *Dynamique et géométrie complexes*, Panoramas et Synthèses **8** (1999)
- [4] R. Kirby and L. Siebenmann, *Foundational essays on topological manifolds, smoothings, and triangulations*, Ann. Math. studies **88** (1977)
- [5] E. Moise, *Geometric topology in dimensions 2 and 3*, Springer–Verlag GTM 47 (1977)
- [6] Y. Rudyak, *Piecewise linear structures on topological manifolds*, eprint arxiv: math.AT/0105047

*Department of Mathematics*  
*Harvard*  
*Cambridge, MA 02138*

Email: [dannyc@math.harvard.edu](mailto:dannyc@math.harvard.edu)

URL: [www.math.harvard.edu/~dannyc](http://www.math.harvard.edu/~dannyc)

Received: 17 May 2001      Revised: 15 August 2001