

Asymptotic Expansions of Integral Functionals of Weakly Correlated Random Processes

J. vom Scheidt, H.-J. Starkloff and R. Wunderlich

Abstract. In the paper asymptotic expansions for second-order moments of integral functionals of a family of random processes are considered. The random processes are assumed to be wide-sense stationary and ε -correlated, i.e. the values are not correlated excluding an ε -neighbourhood of each point. The asymptotic expansions are derived for $\varepsilon \rightarrow 0$. Using a special weak assumption there are found easier expansions as in the case of general weakly correlated random processes. Expansions are given for integral functionals of real-valued as well as of complex vector-valued processes.

Keywords: *Asymptotic expansion, second-order moment, random differential equation, weakly correlated process, stationary process, random vibration*

AMS subject classification: Primary 60 G 12, secondary 34 F 05, 41 A 60, 70 L 05

1. Introduction

In this paper asymptotic expansions for second-order moments of integral functionals of the type

$$\varepsilon_r := \int_{\mathcal{D}} Q(s) \varepsilon f(s) ds \quad (1)$$

are considered, where Q is a deterministic function on an interval $\mathcal{D} \subset \mathbb{R}$ and $(\varepsilon f)_{\varepsilon > 0}$ denotes a family of random functions, indexed by a parameter ε which describes the range of correlation. The random functions are defined on a probability space $(\Omega, \mathfrak{A}, \mathbf{P})$, the expectation operator for random variables on this space is denoted by $\mathbf{E}\{\cdot\}$. Such integral functionals play an important role in many theoretical and practical mathematical problems. For example, differential equations with an inhomogeneous term containing εf often possess solutions which can be represented in such a way (cf. Examples 2 and 3 in this paper). For an approximate description of those solutions and their characteristics asymptotic expansions with respect to $\varepsilon \rightarrow 0$ can be used if the values of εf are correlated or stochastically dependent only in an ε -neighbourhood of each point.

We will suppose the validity of the following

Assumption 1.

1. εf ($\varepsilon > 0$) are wide-sense stationary processes with correlation functions

$$\mathbf{E}\{\varepsilon f(s) \varepsilon f(t)\} = \varepsilon R_{ff}(t - s).$$

All Authors: Techn. Univ. Chemnitz, Fac. Math., D – 09107 Chemnitz

2. ${}^\varepsilon f$ ($\varepsilon > 0$) are centered, i.e. $\mathbf{E}\{ {}^\varepsilon f(s) \} = 0$ for $s \in \mathbb{R}$.
3. ${}^\varepsilon f$ ($\varepsilon > 0$) are ε -correlated, i.e. ${}^\varepsilon R_{ff}(s) = 0$ for $|s| \geq \varepsilon$.
4. The correlation functions ${}^\varepsilon R_{ff}$ ($\varepsilon > 0$) are generated by a correlation function R of a 1-correlated wide-sense stationary process, i.e. ${}^\varepsilon R_{ff}(s) = R(\frac{s}{\varepsilon})$ ($s \in \mathbb{R}$, $\varepsilon > 0$).
5. The correlation function R is continuous on \mathbb{R} , hence the processes ${}^\varepsilon f$ ($\varepsilon > 0$) are continuous in mean square on \mathbb{R} .

The integral in (1) is assumed to exist in mean square sense, under weak conditions it coincides a.s. with the pathwise integral. From condition 2 of Assumption 1 it follows that the random variables ${}^\varepsilon r$ ($\varepsilon > 0$) are centered, i.e. $\mathbf{E}\{ {}^\varepsilon r \} = 0$.

In the first part of the paper we consider real-valued processes, after that complex vector-valued processes are investigated.

For example, $({}^\varepsilon f)_{\varepsilon > 0}$ can be a family of so-called weakly correlated random processes. In the theory of these processes (cf. [8, 9]) asymptotic expansions with respect to $\varepsilon \rightarrow 0$ of the type

$$\mathbf{E}\{ {}^\varepsilon r_1 \cdot {}^\varepsilon r_2 \cdots {}^\varepsilon r_m \} = \begin{cases} c_m \varepsilon^{\frac{m}{2}} + o(\varepsilon^{\frac{m}{2}}) & \text{for even } m \\ c_m \varepsilon^{\frac{m+1}{2}} + o(\varepsilon^{\frac{m+1}{2}}) & \text{for odd } m > 1 \end{cases}$$

with some real constants c_m are derived. The indices 1 to m refer to deterministic functions Q_1, \dots, Q_m and intervals $\mathcal{D}_1, \dots, \mathcal{D}_m$ which are involved in the corresponding integral functionals. Here we will consider only second-order moments and propose a new method of obtaining such asymptotic expansions, which seems to be easier and clarifies in a certain sense the structure of asymptotic expansions in the case of correlation functions. The main difference to the general theory of weakly correlated random processes consists in the explicitly given generating condition 4 of Assumption 1 for the correlation functions ${}^\varepsilon R_{ff}$ ($\varepsilon > 0$).

In the following treatment the concept of correlation moments of wide-sense stationary processes is used.

Definition 1. Let R be a real continuous correlation function of a wide-sense stationary process and $j \in \mathbb{N}_0 = \{0, 1, 2, \dots\}$ with

$$\int_{-\infty}^{\infty} |s|^j |R(s)| ds < \infty.$$

Then

$$\mu_j = \int_{-\infty}^{\infty} s^j R(s) ds = \begin{cases} 0 & \text{for odd } j \\ 2 \int_0^{\infty} s^j R(s) ds & \text{for even } j \end{cases}$$

is called the *correlation moment* of j -th order of the correlation function or the random process and

$$\nu_j = \int_{-\infty}^{\infty} |s|^j R(s) ds = 2 \int_0^{\infty} s^j R(s) ds$$

is called the *absolute correlation moment* of j -th order.

We remark some properties of correlation moments for real-valued wide-sense stationary processes:

1. From the positive definiteness of the correlation function, $\mu_0 = \nu_0 \geq 0$ follows.
2. Property 1 is not true for higher-order correlation moments, i.e. there exist correlation functions and numbers $j \in \mathbb{N}$ with $\nu_j < 0$.
3. For ε -correlated wide-sense stationary processes correlation moments of all orders exist and $\lim_{j \rightarrow \infty} \nu_j = \lim_{j \rightarrow \infty} \mu_j = 0$ holds.

We also note that for 1-correlated wide-sense stationary random processes the following version of the Shannon-Kotelnikov sampling theorem (see, e.g., [3]) is valid:

Proposition 1. *Let*

$$S(\alpha) = \frac{1}{2\pi} \int_{-\infty}^{\infty} R(s) \exp(-i\alpha s) ds = \frac{1}{\pi} \int_0^1 R(s) \cos(\alpha s) ds$$

denote the spectral density of a 1-correlated wide-sense stationary random process. Then the representation

$$S(\alpha) = \lim_{m \rightarrow \infty} \sum_{n=-m}^m \frac{\sin(\alpha - n\pi)}{\alpha - n\pi} S(n\pi)$$

holds for all $\alpha \in \mathbb{R}$, with $\frac{\sin 0}{0} := 1$.

For a non-negative correlation function of a 1-correlated wide-sense stationary random process the corresponding spectral density is a positive definite function. In this case the correlation moments μ_j are closely related to the spectral moments and therefore to the variances of the mean-square derivatives of the adjoint stationary process (cf., e.g., [5: p. 368]).

In the following we will also suppose that the function Q satisfies

Assumption 2. The deterministic function Q is N times continuously differentiable on the interval \mathcal{D} ($N \in \mathbb{N}_0$), $Q^{(N)}$ is absolutely continuous on \mathcal{D} and the derivatives of Q up to the order $N + 1$ belong to the space $L^2(\mathcal{D}) \cap L^1(\mathcal{D})$.

For such functions the Taylor expansion formula with exact integral representation of the remainder (cf., e.g., [4: Section 5.4]) is valid, for the integration by parts formula for absolutely continuous functions see, e.g., [6: Chapter IX/§7].

2. Expansions of variances

From (1) and Assumption 1 it follows that

$$\begin{aligned} \mathbf{E}\{\varepsilon_r^2\} &= \int_{\mathcal{D}} \int_{\mathcal{D}} Q(s)Q(t) \mathbf{E}\{\varepsilon f(s) \varepsilon f(t)\} ds dt \\ &= \int_{\mathcal{D}} \int_{\mathcal{D}} Q(s)Q(t) \varepsilon R_{ff}(t-s) ds dt \\ &= \int_{\mathcal{D}} \int_{\mathcal{D}} Q(s)Q(t) R\left(\frac{t-s}{\varepsilon}\right) ds dt. \end{aligned}$$

The substitution of the variables $t = t$ and $u = \frac{t-s}{\varepsilon}$ gives

$$\mathbf{E}\{\varepsilon r^2\} = \varepsilon \iint_{\varepsilon \mathcal{D}'} Q(t - \varepsilon u)Q(t)R(u) dt du$$

with the transformed domain of integration $\varepsilon \mathcal{D}' = \{(t, u) \in \mathbb{R}^2 : t \in \mathcal{D} \text{ and } t - \varepsilon u \in \mathcal{D}\}$.

In order to show how the method works the case $\mathcal{D} = \mathbb{R}$ is considered explicitly. We deal with the random variables

$$\varepsilon r = \int_{-\infty}^{\infty} Q(s) \varepsilon f(s) ds \quad (\varepsilon > 0)$$

where

$$\begin{aligned} \mathbf{E}\{\varepsilon r^2\} &= \varepsilon \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} Q(t - \varepsilon u)Q(t)R(u) dt du \\ &= \varepsilon \int_{-1}^1 R(u) \int_{-\infty}^{\infty} Q(t - \varepsilon u)Q(t) dt du \end{aligned}$$

can be obtained. Using the notation

$$\phi(u) = \int_{-\infty}^{\infty} Q(t - u)Q(t) dt$$

we write

$$\mathbf{E}\{\varepsilon r^2\} = \varepsilon \int_{-1}^1 \phi(\varepsilon u)R(u) du$$

and it can be seen that the value of the variances depends on the correlation function R and the behavior of the function ϕ in a neighbourhood of zero. Now the Taylor expansion of the function ϕ is applied. From Assumption 2 we find for all $t \in \mathbb{R}$ and $u \in [-1, 1]$

$$Q(t - \varepsilon u) = \sum_{j=0}^N Q^{(j)}(t) \frac{(-\varepsilon u)^j}{j!} + \tilde{\rho}_{N+1}(t, u, \varepsilon)$$

with

$$\tilde{\rho}_{N+1}(t, u, \varepsilon) = \frac{1}{N!} \int_t^{t-\varepsilon u} Q^{(N+1)}(v)(t - \varepsilon u - v)^N dv,$$

and

$$\begin{aligned} \mathbf{E}\{\varepsilon r^2\} &= \sum_{j=0}^N \frac{(-1)^j \varepsilon^{j+1}}{j!} \cdot \int_{-\infty}^{\infty} Q(t)Q^{(j)}(t) dt \cdot \int_{-1}^1 u^j R(u) du \\ &\quad + \frac{\varepsilon}{N!} \int_{-1}^1 \int_{-\infty}^{\infty} \int_t^{t-\varepsilon u} Q^{(N+1)}(v)(t - \varepsilon u - v)^N Q(t)R(u) dv dt du \end{aligned} \tag{2}$$

follows. Integration by parts with respect to the quantities

$$q_j = \int_{-\infty}^{\infty} Q(t)Q^{(j)}(t) dt \quad (j = 0, \dots, N) \tag{3}$$

which do not depend on the random processes leads to

$$\begin{aligned} \int_{-\infty}^{\infty} Q(t)Q^{(j)}(t) dt &= [Q(t)Q^{(j-1)}(t)]_{t \rightarrow -\infty}^{t \rightarrow \infty} - \int_{-\infty}^{\infty} Q'(t)Q^{(j-1)}(t) dt \\ &= - \int_{-\infty}^{\infty} Q'(t)Q^{(j-1)}(t) dt \\ &\quad \vdots \\ &= (-1)^j \int_{-\infty}^{\infty} Q^{(j)}(t)Q(t) dt. \end{aligned}$$

Then

$$\left. \begin{aligned} q_{2k} &= (-1)^k \int_{-\infty}^{\infty} [Q^{(k)}(t)]^2 dt && (k = 0, \dots, [\frac{N}{2}]) \\ q_{2k+1} &= (-1)^{2k+1} q_{2k+1} = 0 && (k = 0, \dots, [\frac{N-1}{2}]) \end{aligned} \right\}$$

follows and the following asymptotic expansion for $\mathbf{E}\{\varepsilon r^2\}$ can be given:

Theorem 1. *Let $(\varepsilon f)_{\varepsilon>0}$ be a family of random processes satisfying Assumption 1 and Q a function satisfying Assumption 2 with $\mathcal{D} = \mathbb{R}$ and $N \in \mathbb{N}_0$. Then*

$$\mathbf{E}\{\varepsilon r^2\} = \sum_{\substack{j=0 \\ j \text{ even}}}^N \frac{\varepsilon^{j+1}}{j!} q_j \mu_j + \rho_{N+1}(\varepsilon),$$

where μ_j denotes the correlation moment of j -th order of the correlation function R , q_j is given in (3) and $\rho_{N+1}(\varepsilon)$ is the last term in (2).

Example 1. For the function $Q(t) = \exp(-\frac{t^2}{2})$,

$$\mathbf{E}\{\varepsilon r^2\} = \sqrt{\pi} \sum_{k=0}^{\infty} \frac{(-1)^k}{4^k k!} \mu_{2k} \varepsilon^{2k+1}$$

holds for all $\varepsilon > 0$. This equation can be proved using the formula

$$Q^{(l)}(t) = (-1)^l H_l(t) \exp\left(-\frac{t^2}{2}\right)$$

where H_l denotes the Hermite polynomial of l -th order with the representation

$$H_{2k}(t) = \sum_{j=0}^k (-1)^j \frac{(2k)!}{2^j j! (2k-2j)!} t^{2(k-j)}$$

for even values $l = 2k$ ($k \in \mathbb{N}$). The convergence to zero of the remainder can be shown using, for example, [1: Estimation 22.14.17].

3. Expansion of correlation functions

Now the mean square continuous wide-sense stationary processes

$$\varepsilon g(t) := \int_{-\infty}^t Q(t-s) \varepsilon f(s) ds = \int_0^{\infty} Q(u) \varepsilon f(t-u) du \quad (\varepsilon > 0)$$

are examined where the deterministic function Q satisfies Assumption 2 on \mathbb{R}_+ . In this case the correlation functions can be written as

$$\begin{aligned} \varepsilon R_{gg}(\tau) &:= \mathbf{E}\{\varepsilon g(t) \varepsilon g(t+\tau)\} \\ &= \int_{-\infty}^t \int_{-\infty}^{t+\tau} Q(t-s_1)Q(t+\tau-s_2) \mathbf{E}\{\varepsilon f(s_1) \varepsilon f(s_2)\} ds_2 ds_1 \\ &= \int_0^{\infty} \int_0^{\infty} Q(u_1)Q(u_2) \varepsilon R_{ff}(\tau+u_1-u_2) du_2 du_1. \end{aligned}$$

Substituting the variables we get

$$\begin{aligned} \varepsilon R_{gg}(\tau) &= \int_{-\infty}^0 \varepsilon R_{ff}(\tau+v_1) \int_{-v_1}^{\infty} Q(v_1+v_2)Q(v_2) dv_2 dv_1 \\ &\quad + \int_0^{\infty} \varepsilon R_{ff}(\tau+v_1) \int_0^{\infty} Q(v_1+v_2)Q(v_2) dv_2 dv_1 \\ &= \int_{-\infty}^0 \varepsilon R_{ff}(\tau+v_1) \int_0^{\infty} Q(v_2)Q(v_2-v_1) dv_2 dv_1 \\ &\quad + \int_0^{\infty} \varepsilon R_{ff}(\tau+v_1) \int_0^{\infty} Q(v_1+v_2)Q(v_2) dv_2 dv_1 \\ &= \int_{-\infty}^{\infty} \varepsilon R_{ff}(\tau+v_1) \int_0^{\infty} Q(v_2+|v_1|)Q(v_2) dv_2 dv_1 \\ &= \int_{-\infty}^{\infty} \varepsilon R_{ff}(w) \int_0^{\infty} Q(u+|w-\tau|)Q(u) du dw \\ &= \int_{-\infty}^{\infty} R\left(\frac{w}{\varepsilon}\right) \int_0^{\infty} Q(u+|w-\tau|)Q(u) du dw \\ &= \varepsilon \int_{-1}^1 R(v) \int_0^{\infty} Q(u+|\varepsilon v-\tau|)Q(u) du dv. \end{aligned}$$

Applying the Taylor expansion of the function $Q(u+|\varepsilon v-\tau|)$ in neighbourhoods of the points $u+|\tau|$,

$$Q(u+|\varepsilon v-\tau|) = \sum_{j=0}^N \frac{1}{j!} Q^{(j)}(u+|\tau|) (|\varepsilon v-\tau| - |\tau|)^j + \tilde{\rho}_{N+1}(u, v, \varepsilon, \tau)$$

follows with a remainder term $\tilde{\rho}_{N+1}(u, v, \varepsilon, \tau)$. Hence

$$\varepsilon R_{gg}(\tau) = \sum_{j=0}^N \frac{\varepsilon}{j!} \int_0^{\infty} Q^{(j)}(u+|\tau|)Q(u) du \int_{-1}^1 R(v) (|\varepsilon v-\tau| - |\tau|)^j dv + \rho_{N+1}(\varepsilon, \tau)$$

with

$$\rho_{N+1}(\varepsilon, \tau) = \varepsilon \int_{-1}^1 R(v) \int_0^\infty \tilde{\rho}_{N+1}(u, v, \varepsilon, \tau) Q(u) du dv$$

holds. Evaluation of the integral terms containing the correlation function R for $\tau \geq 0$ leads to

$$\begin{aligned} & \int_{-1}^1 R(v) (|\varepsilon v - \tau| - |\tau|)^j dv \\ &= \int_{-1}^{\frac{\tau}{\varepsilon} \wedge 1} R(v) (\tau - \varepsilon v - \tau)^j dv + \int_{\frac{\tau}{\varepsilon} \wedge 1}^1 R(v) (\varepsilon v - \tau - \tau)^j dv \\ &= \int_{-1}^{\frac{\tau}{\varepsilon} \wedge 1} R(v) (-\varepsilon v)^j dv + \int_{\frac{\tau}{\varepsilon} \wedge 1}^1 R(v) (\varepsilon v - 2\tau)^j dv \\ &= \int_{-1}^1 R(v) (-\varepsilon v)^j dv + \int_{\frac{\tau}{\varepsilon} \wedge 1}^1 R(v) [(\varepsilon v - 2\tau)^j - (-\varepsilon v)^j] dv \\ &= (-\varepsilon)^j \mu_j + \mathbf{1}_{[0, \varepsilon)}(\tau) \int_{\frac{\tau}{\varepsilon}}^1 R(v) [(\varepsilon v - 2\tau)^j - (-\varepsilon v)^j] dv \end{aligned}$$

where $a \wedge b := \min(a, b)$, and analogously for $\tau \leq 0$ leads to

$$\begin{aligned} & \int_{-1}^1 R(v) (|\varepsilon v - \tau| - |\tau|)^j dv \\ &= \varepsilon^j \mu_j + \mathbf{1}_{(-\varepsilon, 0]}(\tau) \int_{\frac{|\tau|}{\varepsilon}}^1 R(v) [(\varepsilon v - 2|\tau|)^j - (-\varepsilon v)^j] dv. \end{aligned}$$

Using that $\mu_j = 0$ for odd j we have finally:

Theorem 2. *Let $({}^\varepsilon f)_{\varepsilon > 0}$ be a family of random processes satisfying Assumption 1 and Q a function satisfying Assumption 2 with $\mathcal{D} = \mathbb{R}_+$ and $N \in \mathbb{N}_0$. Then*

$${}^\varepsilon R_{gg}(\tau) = \sum_{\substack{j=0 \\ j \text{ even}}}^N \frac{\varepsilon^{j+1}}{j!} q_j(\tau) \mu_j + \mathbf{1}_{(-\varepsilon, \varepsilon)}(\tau) \sum_{j=1}^N \frac{\varepsilon^{j+1}}{j!} q_j(\tau) c_j(\tau) + \rho_{N+1}(\varepsilon, \tau) \quad (4)$$

with the quantities

$$q_j(\tau) = \int_0^\infty Q^{(j)}(u + |\tau|) Q(u) du$$

and

$$c_j(\tau) = \int_{\frac{|\tau|}{\varepsilon}}^1 R(v) \left[\left(v - 2 \frac{|\tau|}{\varepsilon} \right)^j - (-v)^j \right] dv.$$

For fixed values of τ and $\varepsilon \rightarrow 0$ the expansions

$$\left. \begin{aligned} {}^\varepsilon R_{gg}(0) &= \sum_{j=0}^N \frac{\varepsilon^{j+1}}{j!} q_j(0) \nu_j + o(\varepsilon^{N+1}) \\ {}^\varepsilon R_{gg}(\tau) &= \sum_{\substack{j=0 \\ j \text{ even}}}^N \frac{\varepsilon^{j+1}}{j!} q_j(\tau) \mu_j + o(\varepsilon^{N+1}) \quad (\tau \neq 0) \end{aligned} \right\}$$

are valid. We can see that a discontinuity in the expansions of the correlation function at the point $\tau = 0$ arises if $q_j(0)$ or ν_j for odd values j do not vanish.

Examining asymptotic expansions of ${}^\varepsilon R_{gg}(\tau)$ as a function of τ it is necessary to consider not only the first terms in (4) but also the correction terms in the second sum of (4) for $|\tau| < \varepsilon$.

Example 2. The stationary solution of the second-order linear differential equation with constant coefficients and a random weakly correlated wide-sense stationary inhomogeneous term

$$\ddot{x} + 2\delta\dot{x} + w_0^2x = {}^\varepsilon f(t)$$

is given by

$$x(t) = \frac{1}{w} \int_{-\infty}^t e^{-\delta(t-s)} \sin(w(t-s)) {}^\varepsilon f(s) ds$$

with $w = \sqrt{w_0^2 - \delta^2}$, $0 < \delta < w_0$. In this case the kernel function of the integral functional reads as

$$Q(u) = \frac{1}{w} e^{-\delta u} \sin(wu)$$

and straightforward calculations lead to

$$q_j(\tau) = a_j e^{-\delta|\tau|} \cos(w|\tau|) + b_j e^{-\delta|\tau|} \sin(w|\tau|) \quad (j \in \mathbb{N}, \tau \in \mathbb{R})$$

with

$$\left. \begin{aligned} a_j &= \sum_{l=0}^j \binom{j}{l} (-\delta)^l w^{j-l-2} \int_0^\infty e^{-2\delta u} \sin\left(wu + (j-l)\frac{\pi}{2}\right) \sin(wu) du \\ b_j &= \sum_{l=0}^j \binom{j}{l} (-\delta)^l w^{j-l-2} \int_0^\infty e^{-2\delta u} \cos\left(wu + (j-l)\frac{\pi}{2}\right) \sin(wu) du. \end{aligned} \right\}$$

For example, from the relations

$$\left. \begin{aligned} \int_0^\infty e^{-2\delta u} \sin^2(wu) du &= \frac{1}{4} \frac{w^2}{\delta(\delta^2 + w^2)} \\ \int_0^\infty e^{-2\delta u} \cos(wu) \sin(wu) du &= \frac{1}{4} \frac{w}{\delta^2 + w^2}, \end{aligned} \right\}$$

for $j = 0, 1, 2$ and $\tau \in \mathbb{R}$

$$\left. \begin{aligned} q_0(\tau) &= \frac{e^{-\delta|\tau|}}{4(\delta^2 + w^2)} \left(\frac{1}{\delta} \cos(\omega|\tau|) + \frac{1}{w} \sin(\omega|\tau|) \right) \\ q_1(\tau) &= -\frac{e^{-\delta|\tau|}}{4\delta w} \sin(\omega|\tau|) \\ q_2(\tau) &= \frac{e^{-\delta|\tau|}}{4} \left(-\frac{1}{\delta} \cos(\omega|\tau|) + \frac{1}{w} \sin(\omega|\tau|) \right) \end{aligned} \right\}$$

follows. Choosing the hat-like correlation function

$$R(v) = \begin{cases} 1 - |v| & \text{for } |v| \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

the correlation moments can be found by

$$\mu_j = \frac{2}{(j+1)(j+2)}$$

for even values of j , and for the correction terms we get by setting $\alpha := \frac{|\tau|}{\varepsilon}$

$$\begin{aligned} c_j(\tau) &= \int_{\alpha}^1 R(v) [(v - 2\alpha)^j - (-v)^j] dv \\ &= \frac{2(-1)^{j+1}}{j+1} (\alpha - 1)\alpha^{j+1} + \frac{(1 - 2\alpha)^{j+2} - (-1)^j}{(j+1)(j+2)}. \end{aligned}$$

Especially,

$$\left. \begin{aligned} c_1(\tau) &= \frac{1}{3} - \alpha + \alpha^2 - \frac{1}{3}\alpha^3 \\ c_2(\tau) &= -\frac{2}{3}\alpha + 2\alpha^2 - 2\alpha^3 + \frac{2}{3}\alpha^4. \end{aligned} \right\}$$

4. Expansion of covariance matrices

Now complex vector-valued processes are investigated. In this case the non-commutativity of matrix multiplication has to be taken into account.

We assume the validity of the corresponding versions of Assumption 1 for \mathbb{C}^n -valued wide-sense stationary processes ${}^\varepsilon f$ ($\varepsilon > 0$), and of Assumption 2 for $m \times n$ -matrix-valued deterministic functions Q ($m, n \in \mathbb{N}$). So for the matrix correlation functions of the \mathbb{C}^n -valued wide-sense stationary processes ${}^\varepsilon f$ ($\varepsilon > 0$)

$${}^\varepsilon R_{ff}(s) := \mathbf{E}\{ {}^\varepsilon f(t) {}^\varepsilon f^*(t+s) \} = R\left(\frac{s}{\varepsilon}\right)$$

holds according to condition 4 of Assumption 1. Here $*$ denotes the conjugate-complex transposed of matrices and vectors where vectors are assumed to be columns. From the relation $R(s) = R^*(-s)$ we find in this case for the correlation moments

$$\left. \begin{aligned} \mu_j &= \int_{-\infty}^{\infty} s^j R(s) ds = \nu_j^+ + (-1)^j [\nu_j^+]^* \\ \nu_j &= \int_{-\infty}^{\infty} |s|^j R(s) ds = \nu_j^+ + [\nu_j^+]^* \end{aligned} \right\}$$

with $\nu_j^+ = \int_0^{\infty} s^j R(s) ds$, hence $\mu_j^* = (-1)^j \mu_j$ and $\nu_j^* = \nu_j$ hold. For functionals of the type

$${}^\varepsilon r = \int_{-\infty}^{\infty} Q(s) {}^\varepsilon f(s) ds \quad (\varepsilon > 0)$$

we get

$$\mathbf{E}\{ {}^\varepsilon r {}^\varepsilon r^* \} = \varepsilon \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} Q(t - \varepsilon u) R(u) Q^*(t) dt du$$

and we find in the same way as for real-valued processes:

Theorem 3. *Let $(\varepsilon f)_{\varepsilon>0}$ be a family of ε -correlated \mathbb{C}^n -valued wide-sense stationary random processes satisfying Assumption 1 and Q a matrix-valued function satisfying Assumption 2 on $\mathcal{D} = \mathbb{R}$ and $N \in \mathbb{N}_0$. Then*

$$\mathbf{E}\{\varepsilon_r \varepsilon_r^*\} = \sum_{j=0}^N \frac{\varepsilon^{j+1}}{j!} q_j + \rho_{N+1}(\varepsilon)$$

with

$$q_j = \int_{-\infty}^{\infty} Q^{(j)}(t) \mu_j^* Q^*(t) dt$$

and the remainder term

$$\rho_{N+1}(\varepsilon) = \frac{\varepsilon}{N!} \int_{-1}^1 \int_{-\infty}^{\infty} \int_t^{t-\varepsilon u} Q^{(N+1)}(v) (t - \varepsilon u - v)^N R(u) Q^*(t) dv dt du.$$

5. Expansion of matrix correlation functions

Considering the complex vector-valued wide-sense stationary processes

$$\varepsilon g(t) := \int_{-\infty}^t Q(t-s) \varepsilon f(s) ds = \int_0^{\infty} Q(u) \varepsilon f(t-u) du \quad (\varepsilon > 0)$$

the matrix correlation functions can be written as

$$\varepsilon R_{gg}(\tau) := \mathbf{E}\{\varepsilon g(t) \varepsilon g^*(t+\tau)\} = \int_0^{\infty} \int_0^{\infty} Q(u_1) \varepsilon R_{ff}(\tau + u_1 - u_2) Q^*(u_2) du_2 du_1$$

and it can be obtained

$$\begin{aligned} \varepsilon R_{gg}(\tau) &= \int_{-\infty}^0 \int_{-v_1}^{\infty} Q(v_1 + v_2) \varepsilon R_{ff}(\tau + v_1) Q^*(v_2) dv_2 dv_1 \\ &\quad + \int_0^{\infty} \int_0^{\infty} Q(v_1 + v_2) \varepsilon R_{ff}(\tau + v_1) Q^*(v_2) dv_2 dv_1 \\ &= \int_{-\infty}^0 \int_0^{\infty} Q(v_2) \varepsilon R_{ff}(\tau + v_1) Q^*(v_2 - v_1) dv_2 dv_1 \\ &\quad + \int_0^{\infty} \int_0^{\infty} Q(v_1 + v_2) \varepsilon R_{ff}(\tau + v_1) Q^*(v_2) dv_2 dv_1 \\ &= \int_{-\infty}^{\infty} \int_0^{\infty} Q(v_2 + v_1^+) \varepsilon R_{ff}(\tau + v_1) Q^*(v_2 + v_1^-) dv_2 dv_1 \\ &= \int_{-\infty}^{\infty} \int_0^{\infty} Q(u + (w - \tau)^+) \varepsilon R_{ff}(w) Q^*(u + (w - \tau)^-) dudw \\ &= \varepsilon \int_{-1}^1 \int_0^{\infty} Q(u + (\varepsilon v - \tau)^+) R(v) Q^*(u + (\varepsilon v - \tau)^-) dudv. \end{aligned}$$

Now, Taylor expansion of the matrix function Q in neighbourhoods of the points $u + |\tau|$ is considered. In the case of $\tau = 0$,

$$\begin{aligned} \varepsilon R_{gg}(0) &= \varepsilon \int_0^1 \int_0^\infty Q(u + |\varepsilon v|) R(v) Q^*(u) \, dudv \\ &\quad + \varepsilon \int_{-1}^0 \int_0^\infty Q(u) R(v) Q^*(u + |\varepsilon v|) \, dudv \\ &= \sum_{j=0}^N \frac{\varepsilon^{j+1}}{j!} \left\{ \int_0^1 \int_0^\infty Q^{(j)}(u) v^j R(v) Q^*(u) \, dudv \right. \\ &\quad \left. + \left[\int_0^1 \int_0^\infty Q^{(j)}(u) v^j R(v) Q^*(u) \, dudv \right]^* \right\} + \rho_{N+1}(\varepsilon, 0) \\ &= \sum_{j=0}^N \frac{\varepsilon^{j+1}}{j!} \{q_j + q_j^*\} + \rho_{N+1}(\varepsilon, 0) \end{aligned}$$

with

$$q_j = \int_0^\infty Q^{(j)}(u) \nu_j^+ Q^*(u) \, du.$$

In an analogous manner to Section 3 we obtain for $\tau > 0$

$$\begin{aligned} \varepsilon R_{gg}(\tau) &= \sum_{j=0}^N \frac{\varepsilon^{j+1}}{j!} \left\{ q_j(\tau) \right. \\ &\quad + \mathbf{1}_{(0,\varepsilon)}(\tau) \left\{ \int_0^\infty Q^{(j)}(u + \tau) \int_{\frac{\tau}{\varepsilon}}^1 \left(v - \frac{2\tau}{\varepsilon}\right)^j R(v) \, dv Q^*(u) \, du \right. \\ &\quad \left. \left. + (-1)^{j+1} \int_0^\infty Q(u) \int_{\frac{\tau}{\varepsilon}}^1 v^j R(v) \, dv [Q^{(j)}(u + \tau)]^* \, du \right\} \right\} + \rho_{N+1}(\varepsilon, \tau) \end{aligned}$$

with

$$q_j(\tau) = \int_0^\infty Q(u) \mu_j^* [Q^{(j)}(u + \tau)]^* \, du$$

and the following theorem holds:

Theorem 4. *Let $(\varepsilon f)_{\varepsilon>0}$ be a family of \mathbb{C}^n -valued random processes satisfying Assumption 1 and Q a matrix-valued function satisfying Assumption 2 on $\mathcal{D} = \mathbb{R}_+$ and $N \in \mathbb{N}_0$. Then for $\tau > 0$*

$$\varepsilon R_{gg}(\tau) = \sum_{j=0}^N \frac{\varepsilon^{j+1}}{j!} \{q_j(\tau) + \mathbf{1}_{(0,\varepsilon)}(\tau) c_j(\tau)\} + \rho_{N+1}(\varepsilon, \tau)$$

with

$$q_j(\tau) = \int_0^\infty Q(u) \mu_j^* [Q^{(j)}(u + \tau)]^* \, du$$

and

$$c_j(\tau) = \int_0^\infty Q^{(j)}(u + \tau) \int_{\frac{\tau}{\varepsilon}}^1 \left(v - \frac{2\tau}{\varepsilon}\right)^j R(v) dv Q^*(u) du + (-1)^{j+1} \int_0^\infty Q(u) \int_{\frac{\tau}{\varepsilon}}^1 v^j R(v) dv [Q^{(j)}(u + \tau)]^* du.$$

For $\tau = 0$, the relation

$${}^\varepsilon R_{gg}(0) = \sum_{j=0}^N \frac{\varepsilon^{j+1}}{j!} \{q_j + q_j^*\} + \rho_{N+1}(\varepsilon, 0)$$

is valid with

$$q_j = \int_0^\infty Q^{(j)}(u) \nu_j^+ Q^*(u) du$$

and the values for $\tau < 0$ can be calculated from the relation ${}^\varepsilon R_{gg}(\tau) = {}^\varepsilon R_{gg}^*(-\tau)$.

Example 3. Let us consider a system of n ordinary differential equations of first order with constant coefficients and an ε -correlated wide-sense stationary inhomogeneous term

$${}^\varepsilon \dot{x} = A {}^\varepsilon x + {}^\varepsilon f.$$

The matrix A is assumed to be stable (i.e. all eigenvalues $\lambda_1, \dots, \lambda_n$ have negative real parts) and diagonalizable (i.e. $A = V \Lambda V^{-1}$ with $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$). Then a stationary solution

$${}^\varepsilon x(t) = \int_{-\infty}^t e^{A(t-s)} {}^\varepsilon f(s) ds$$

of the system exists (see, e.g., [2, 7, 9]).

Then for $\tau = 0$

$${}^\varepsilon R_{xx}(0) = \sum_{j=0}^N \frac{\varepsilon^{j+1}}{j!} \{q_j + q_j^*\} + o(\varepsilon^{N+1})$$

as $\varepsilon \rightarrow 0$ with

$$q_j = V \Lambda^j \int_0^\infty e^{\Lambda u} B_j e^{\Lambda^* u} du V^* = -V \left(\frac{b_{jkl} \lambda_k^j}{\lambda_k + \bar{\lambda}_l} \right)_{k,l=1}^n V^* \\ B_j = (b_{jkl})_{k,l=1}^n = V^{-1} \nu_j^+ [V^{-1}]^*$$

and for fixed $\tau > 0$

$${}^\varepsilon R_{xx}(\tau) = \sum_{j=0}^N \frac{\varepsilon^{j+1}}{j!} q_j(\tau) + o(\varepsilon^{N+1})$$

as $\varepsilon \rightarrow 0$ with

$$q_j(\tau) = V \int_0^\infty e^{\Lambda u} C_j e^{\Lambda^* u} du [\Lambda^j]^* e^{\Lambda^* \tau} V^* = -V \left(\frac{c_{jkl} e^{\bar{\lambda}_l \tau} \bar{\lambda}_l^{-j}}{\lambda_k + \bar{\lambda}_l} \right)_{k,l=1}^n V^* \\ C_j = (c_{jkl})_{k,l=1}^n = V^{-1} \mu_j^* [V^{-1}]^*$$

holds.

6. Conclusion

Asymptotic expansions as $\varepsilon \rightarrow 0$ of variances or correlation functions of integral functionals involving ε -correlated wide-sense stationary random functions have been derived. In these expansions the influence of the deterministic kernel function and of the random function can be separated using the concept of correlation moments and certain characteristics of the kernel function. In the case of random variables the expansions have the form of a power series in ε , in the case of a correlation function additional correction terms for $|\tau| < \varepsilon$ arise. For given kernel functions and a generating correlation function it is easy to compute (at least numerically) the terms of the expansion. An estimation of the remainder term is also possible.

With respect to applications it is worth to note that the asymptotic expansions can over- or underestimate the true value. The statement of the overestimation of the true value in [9: p. 49] results from the special type of the correlation functions considered there.

Further expansions of second-order characteristics of integral functionals of random processes can be found in [10, 11]. An extension of the results to scalar- or vector-valued random fields and certain classes of non-stationary processes is also possible and will be considered in a subsequent paper.

Acknowledgment. The support by the Deutsche Forschungsgemeinschaft is gratefully appreciated.

References

- [1] Abramowitz, M. and I. A. Stegun: *Pocketbook of Mathematical Functions*. Thun, Frankfurt/Main: Verlag Harri Deutsch 1984.
- [2] Bunke, H.: *Gewöhnliche Differentialgleichungen mit zufälligen Parametern*. Berlin: Akademie-Verlag 1972.
- [3] Butzer, P. L., Splettstößer, W. and R. L. Stens: *The sampling theorem and linear prediction in signal analysis*. Jahresber. Deutsch. Math.-Verein. 90 (1988), 1 – 70.
- [4] Courant, R. and F. John: *Introduction to Calculus and Analysis*, Vol. I. New York: Springer-Verlag 1989.
- [5] Gnedenko, B. W.: *Einführung in die Wahrscheinlichkeitstheorie*. Berlin: Akademie-Verlag 1991.
- [6] Natanson, I. P.: *Theorie der Funktionen einer reellen Veränderlichen*. Berlin: Akademie-Verlag 1975.
- [7] Preumont, A.: *Random Vibration and Spectral Analysis*. Dordrecht: Kluwer 1994.
- [8] Scheidt, J. vom: *Stochastic Equations of Mathematical Physics*. Berlin: Akademie-Verlag 1990.
- [9] Scheidt, J. vom, Fellenberg, B. and U. Wöhl: *Analyse und Simulation stochastischer Schwingungssysteme*. Stuttgart: B. G. Teubner 1994.
- [10] Scheidt, J. vom, Starkloff, H.-J. and R. Wunderlich: *Asymptotic expansions for second-order moments of integral functionals of weakly correlated random functions*. Preprint. Chemnitz: Fac. Math. Techn. Univ., Preprint 97-17 (1997), pp. 1 – 9.

http://www.mathematik.tu-chemnitz.de/preprint/1997/PREPRINT_17.html.

- [11] Scheidt, J. vom, Starkloff, H.-J. and R. Wunderlich: *Asymptotic expansions of integral functionals of vector valued ε -correlated processes*. Preprint. Chemnitz: Fac. Math. Techn. Univ., Preprint 99-03 (1999), pp. 1 – 13.

http://www.mathematik.tu-chemnitz.de/preprint/1999/PREPRINT_03.html.

Received 20.04.1999